

527 rec'd PCT/EP

06 NOV 2000

TRANSMIT A LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

0147-0215P

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

NEW

INTERNATIONAL APPLICATION NO.

PCT/EP99/03292

INTERNATIONAL FILING DATE

May 12, 1999

PRIORITY DATE CLAIMED

09/674768
May 13, 1998

TITLE OF INVENTION

TRANSGENIC PLANTS WITH A MODIFIED ACTIVITY OF A PLASTIDIAL ADP/ATP TRANSLOCATOR

APPLICANT(S) FOR DO/EO/US

NEUHAUS, Ekkehard; MOEHLMANN, Torsten; GRAEVE-KAMPFENKEL, Karl-Heinz; TJADEN, Joachim *

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39 (1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☒ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☒ A translation of the International Application into English (35 U.S.C. 371(c)(3)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(2)).
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern document(s) or information included:

11. ☒ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.-1449 and International Search Report (PCT/ISA/210 in German) w/ 8 documents
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ A FIRST preliminary amendment.
☐ A SECOND or SUBSEQUENT preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information:
 - 1.) Article 34 amended claims
 - 2.) Eight (8) sheets of Formal Drawings

526 Rec'd PCT/PTO 06 NOV 2000

U.S. APPLICATION NO. (if known, see 37 CFR 1.5) 09/ 674768		INTERNATIONAL APPLICATION NO. PCT/EPP99/03292		ATTORNEY'S DOCKET NUMBER 0147-0215P	
--	--	--	--	--	--

17. <input checked="" type="checkbox"/> The following fees are submitted: BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5): Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO. \$1,000.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$860.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO. \$710.00 International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$690.00 International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) \$100.00 ENTER APPROPRIATE BASIC FEE AMOUNT =	CALCULATIONS PTO USE ONLY																					
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input checked="" type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).	\$ 860.00																					
<table border="1" style="width:100%; border-collapse: collapse;"> <tr> <th style="width:20%;">CLAIMS</th> <th style="width:20%;">NUMBER FILED</th> <th style="width:20%;">NUMBER EXTRA</th> <th style="width:20%;">RATE</th> </tr> <tr> <td>Total Claims</td> <td>25 - 20 =</td> <td>5</td> <td>X \$18.00</td> </tr> <tr> <td>Independent Claims</td> <td>3 - 3 =</td> <td>0</td> <td>X \$80.00</td> </tr> <tr> <td colspan="3">MULTIPLE DEPENDENT CLAIM(S) (if applicable) Yes</td> <td>+ \$270.00</td> </tr> <tr> <td colspan="3">TOTAL OF ABOVE CALCULATIONS =</td> <td>\$ 1350.00</td> </tr> </table>	CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	Total Claims	25 - 20 =	5	X \$18.00	Independent Claims	3 - 3 =	0	X \$80.00	MULTIPLE DEPENDENT CLAIM(S) (if applicable) Yes			+ \$270.00	TOTAL OF ABOVE CALCULATIONS =			\$ 1350.00	\$ 130.00	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE																			
Total Claims	25 - 20 =	5	X \$18.00																			
Independent Claims	3 - 3 =	0	X \$80.00																			
MULTIPLE DEPENDENT CLAIM(S) (if applicable) Yes			+ \$270.00																			
TOTAL OF ABOVE CALCULATIONS =			\$ 1350.00																			
Reduction of 1/2 for filing by small entity, if applicable. Verified Small Entity statement must also be filed (Note 37 CFR 1.9, 1.27, 1.28).	\$ 0																					
SUBTOTAL =	\$ 1350.00																					
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).	\$ 0																					
TOTAL NATIONAL FEE =	\$ 1350.00																					
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property +	\$ 0																					
TOTAL FEES ENCLOSED =	\$ 1350.00																					
<table border="1" style="width:100%; border-collapse: collapse;"> <tr> <td style="width:60%;"></td> <td style="width:20%; text-align: center;">Amount to be:</td> <td style="width:20%; text-align: center;">\$</td> </tr> <tr> <td></td> <td style="text-align: center;">refunded</td> <td></td> </tr> <tr> <td></td> <td style="text-align: center;">charged</td> <td style="text-align: center;">\$</td> </tr> </table>				Amount to be:	\$		refunded			charged	\$											
	Amount to be:	\$																				
	refunded																					
	charged	\$																				


a. ☒ A check in the amount of \$ **1350.00** to cover the above fees is enclosed.

b. ☐ Please charge my Deposit Account. No. _____ in the amount of \$ _____ to cover the above fees.
 A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any
 overpayment to Deposit Account No. 02-2448.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR
 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

Send all correspondence to:
Birch, Stewart, Kolasch & Birch, LLP or Customer No. 2292
 P.O. Box 747
 Falls Church, VA 22040-0747
 (703)205-8000


 SIGNATURE

SVENSSON, LEONARD R.
 NAME

#30,330 (LRS)
 REGISTRATION NO.

#3/a

IN THE U.S. PATENT AND TRADEMARK OFFICE

Applicant: NEUHAUS, Ekkehard et al.
Int'l. Appl. No.: PCT/EP99/03292
Appl. No.: New Group:
Filed: November 6, 2000 Examiner:
For: TRANSGENIC PLANTS WITH A MODIFIED
ACTIVITY OF A PLASTIDIAL ADP/ATP
TRANSLOCATOR

PRELIMINARY AMENDMENT

BOX PATENT APPLICATION

Assistant Commissioner for Patents
Washington, DC 20231

November 6, 2000

Sir:

The following Preliminary Amendments and Remarks are respectfully submitted in connection with the above-identified application.

AMENDMENTS

IN THE SPECIFICATION:

Please amend the specification as follows:

Before line 1, insert --This application is the national phase under 35 U.S.C. § 371 of PCT International Application No. PCT/EP99/03292 which has an International filing date of May 12, 1999, which designated the United States of America.--

Page 26: Line 24, after "AATP2" insert --(SEQ ID NO:6)--

Line 25, after "(A.thaliana" insert --SEQ ID NO:5--;and

Line 26, after "Rickettsia prowasekii" insert --(SEQ ID NO:7)--.

Docket No. 0147-0215P

IN THE CLAIMS:

Please amend the claims as follows:

Claim 5: Line 1, change "any one of claims 1 to 4" to --
claim 1--

Claim 6: Line 1, change "any one of claims 1 to 5" to --
claim 1--

Claim 7: Lines 1 to 2, change "any one of claims 1 to 6"
to --claim 1--

Claim 13: Lines 2 to 3, change "any one of claims 1 to 6"
to --claim 1--

Claim 16: Lines 2 to 3, delete "or according to claim 12"

REMARKS

The specification has been amended to provide a cross-reference to the previously filed International Application. The claims have also been amended to delete improper multiple dependencies and to place the application into better form for examination.

Enclosed herewith in full compliance with 37 C.F.R. §§ 1.821-1825 is a Substitute Sequence Listing to be inserted into the specification as indicated above. the Substitute Sequence Listing in no way introduces new matter into the specification. Also submitted herewith in full compliance with 37 C.F.R. §§ 1.821-1825 is a computer disk containing the computer readable form of the Substitute Sequence Listing, file "0147-215.txt", is identical to the paper copy, except that it lacks formatting.

Docket No. 0147-0215P

Entry of the present amendment and favorable action on the above-identified application are respectfully requested.

If necessary, the Commissioner is hereby authorized in this, concurrent, and future replies, to charge payment or credit any overpayment to Deposit Account No. 02-2448 for any additional fees required under 37 C.F.R. § 1.16 or under 37 C.F.R. § 1.17; particularly, extension of time fees.

Respectfully submitted,

BIRCH, STEWART, KOLASCH & BIRCH, LLP

By  #32,868
for Leonard R. Svensson, #30,330

LRS/lps:cgc
0147-0215P

P.O. Box 747
Falls Church, VA 22040-0747
(703) 205-8000

(Rev. 04/19/2000)

8/PR+5

526 Rec'd PCT/PTO 06 NOV 2000

VOSSIUS & PARTNER

Patentanwälte

09/ 674768

SIEBERTSTRASSE 4 · 81675 MÜNCHEN

TEL.: +49-89-4130 40 · FAX: +49-89-4130 41 11 · FAX (Marken-Trademarks): +49-89-41 30 44 00

5

Our Ref.: C 1540 PCT

10

**Transgenic plants with modified activity of
a plastidial ADP/ADP translocator**

15

The present invention relates to transgenic plant cells and plants with an increased plastidial ADP/ATP translocator activity. Such cells and plants exhibit an increased yield, preferably an increased oil and/or starch content, and synthesize preferably a starch with increased amylose content.

Further, the present invention relates to transgenic plant cells and plants with a decreased ADP/ATP translocator activity. Such cells and plants synthesize a starch with decreased amylose content.

20

In the field of agriculture and forestry there have been permanent endeavours to provide plants with an increased yield, in particular, in order to ensure the food supply of the permanently growing world population and to guarantee the supply of regenerating raw materials. Traditionally, it has been tried to obtain high-yield plants by breeding. This, however, is time-consuming and costly. Furthermore, corresponding breeding programs have to be carried out for each plant species of interest.

25

Progress has been made, partly, by genetic manipulation of plants, i.e. by purposeful introduction and expression of recombinant nucleic acid molecules in plants.

30

Approaches of that kind have the advantage that they are, in general, not limited to one plant species but can be transferred to other plant species, too.

35

In EP-A 0 511 979, for example, it was described that the expression of a prokaryotic asparagin synthetase in plant cells leads to an increased biomass production, amongst others. WO 96/21737 describes, for example, the increase in yield of plants by expression of de- or unregulated fructose-1,6-bisphosphatase due to increase in the rate of photosynthesis. Nevertheless, there is still a need for generally applicable methods for the improvement of the yield in plants interesting for agriculture or forestry.

40

Furthermore, with regard to the fact that substances contained in plants play a more and more important role as renewable sources of raw material, one of the problems in biotechnological research is the adjustment of said vegetable raw materials to the requirements of the processing industry. In order to allow for application of

regenerating raw materials in as many fields as possible it is furthermore necessary to achieve a wide range of substances. Moreover, it is necessary to increase the yield of said vegetable content substances in order to increase the efficiency of the production of renewable sources of raw material from plants.

- 5 Apart from fats and proteins, oils and polysaccharides are the essential regenerating vegetable raw materials. A central role with the polysaccharides, apart from cellulose, plays starch which is one of the most important reserve substances in higher plants. Amongst those, potato and maize, in particular, are interesting plants since they are important cultivated plants for the production of starch.
- 10 The polysaccharide starch which is one of the most important reserve substances in the vegetable world is, apart from its use in the food industry, widely used as regenerating raw material for the production of industrial products.

- 15 The starch industry has a great interest in plants with increased starch content which, as a rule, means an increased dry weight. An increased dry weight increases the value of the plants processed in the starch industry (maize, potato, tapioca, wheat, barley, rice etc.) due to the increased yield of starch. In addition, plant cells or organs containing higher amounts of starch offer advantages for the processing in the food industry since they absorb less fat or frying oil and, thus, lead to "healthier" products.
- 20 with reduced caloric content. Said property is of great importance e.g. in the production of popcorn, corn flakes from maize or chips, crisps or potato fritters from potatoes.

- 25 For the industry processing potato starch the dry weight (starch content) is a crucial size since it determines processing costs. An increased dry weight (starch content) means, that with the same yield, the water content of the potato tuber is reduced. The reduced water content leads to reduced transport costs and to a reduction of the exact cooking period necessary in cooking.

- 30 Therefore, it seems desirable to provide plant cells and plants exhibiting an increased starch content as well as methods for the production of such plant cells and plants. Moreover, it seems desirable to provide starches whose amylose and amylopectin content meets the requirements of the processing industry. In this context, both starches with an increased amylose content and starches with a reduced amylose content are of interest since they are particularly suitable for special uses each.

35

Thus, the problem underlying the present invention is to provide plant cells and plants which, in comparison with corresponding non-modified wild type plant cells and wild type plants, exhibit an increased yield preferably of oil and/or starch and/or synthesize a starch with a modified amylose content.

This problem is solved by the provision of the embodiments characterized in the claims.

- 5 Thus, the present invention relates to transgenic plant cells which are genetically modified, wherein the genetic modification is the introduction of a foreign nucleic acid molecule whose presence or expression leads to an increase in the plastidial ADP/ATP translocator activity in the transgenic cells in comparison with corresponding genetically non-modified plant cells from wild type plants.

10

- In this context, the genetic modification can be any genetic modification leading to an increase in the plastidial ADP/ATP translocator activity. One possibility, for example, is the so-called "in situ-activation", wherein the genetic modification is a change of the regulatory regions of endogenous ADP/ATP translocator genes, which leads to an increased expression of said genes. This can be achieved, for example, by means of introduction of a very strong promoter in front of the corresponding genes, e.g. by means of homologous recombination.

15

- Further, there is the possibility to apply the method of the so-called "activation tagging" (cf. e.g. Walden et al., Plant J. (1991), 281-288; Walden et al., Plant Mol. Biol. 26 (1994), 1521-1528). Said method is based on the activation of endogenous promoters by means of enhancer elements such as the enhancer of the 35S RNA promoter of the cauliflower mosaic virus or the octopin synthase enhancer.

20

- 25 In a preferred embodiment the genetic modification comprises, however, the introduction of a foreign nucleic acid molecule encoding a plastidial ADP/ATP translocator into the genome of the plant cell.

- The term "transgenic", therefore, means that the plant cell of the invention contains at least one foreign nucleic acid molecule encoding a plastidial ADP/ATP translocator stably integrated in the genome, preferably a nucleic acid molecule.

30

- The term "foreign nucleic acid molecule" preferably means a nucleic acid molecule which encodes a protein with the biological activity of a plastidial ADP/ATP translocator and either does not occur naturally in corresponding plant cells or does not occur naturally in the precise spatial order in the plant cells or which is localized at a place in the genome of the plant cell where it does not occur naturally. Preferably, the foreign nucleic acid molecule is a recombinant molecule which consists of various elements and whose combination or specific spatial arrangement

35

does not occur naturally in plant cells. The transgenic plant cells of the invention contain at least one foreign nucleic acid molecule encoding a protein with the biological activity of a plastidial ADP/ATP translocator, wherein said nucleic acid molecule preferably is connected with regulatory DNA elements ensuring the transcription in plant cells, in particular with a promoter.

In principle, the foreign nucleic acid molecule can be any nucleic acid molecule encoding an ADP/ATP translocator which, after expression, is localized in the inner membrane of plastids. In this context, a plastidial ADP/ATP translocator is a protein catalyzing the transport of ATP into the plastids and of ADP out of the plastids. Such nucleic acid molecules are known, for example, from *Arabidopsis thaliana* (Kampfenkel et al., FEBS Lett. 374 (1995), 351-355; Genbank Acc. No. X94626 and Acc. No. Z49227) or from potato (Genbank Acc. No. Y10821). By means of said known nucleic acid molecules the person skilled in the art can isolate corresponding sequences from other organisms, particularly vegetable ones, according to standard methods, for example by heterologous screening. In particular, non-vegetable nucleic acid molecules can be used, too, which encode an ADP/ATP translocator and are connected with a targeting sequence ensuring the localisation in the inner plastid membrane. In this context, e.g. an ADP/ATP translocator is known from *Rickettsia prowazekii* (Williamson et al., Gene 80 (1989), 269-278) and from *Chlamydia trachomatis*.

In a preferred embodiment the foreign nucleic acid molecule encodes a plastidial ADP/ATP translocator from *Arabidopsis thaliana*, in particular the protein AATP1 described in Kampfenkel et al. (1995, loc. cit.).

The cells of the invention can be distinguished from naturally occurring plant cells, amongst others, in that they contain a foreign nucleic acid molecule which does not occur naturally in said cells or in that said molecule is integrated at a place in the genome of the cell where it does not normally occur, i.e. in another genomic environment. Further, said transgenic plant cells of the invention can be differentiated from naturally occurring plant cells in that they contain at least one copy of the foreign nucleic acid molecule stably integrated in their genome, optionally in addition to the copies of said molecule naturally occurring in the cells. If the nucleic acid molecule(s) introduced into the cells is/ (are) an additional copy(ies) of molecules naturally occurring in the cells, the plant cells of the invention can be differentiated from naturally occurring plant cells particularly in that said additional copy(ies) is/ (are) located at places in the genome where it (they) (do)/ does not occur naturally. This can, for example, be determined by means of a Southern blot analysis.

The plant cells of the invention can further be differentiated from naturally occurring plant cells preferably by at least one of the following features: If the nucleic acid molecule is heterologous regarding the plant cell, the transgenic plant cells exhibit transcripts of the nucleic acid molecule introduced, which can be detected by e.g. Northern blot analysis. Preferably, the plant cells of the invention contain a protein which is encoded by a nucleic acid molecule introduced. This can be detected by e.g. immunological methods, particularly by Western blot analysis.

If the nucleic acid molecule is homologue regarding the plant cell, the cells of the invention can be differentiated from naturally occurring cells, for example, due to the additional expression of the foreign nucleic acid molecules introduced. Preferably, the transgenic plant cells contain more transcripts of the foreign nucleic acid molecules. This can be detected by e.g. Northern blot analysis.

The term "genetically modified" means that the plant cell is modified in its genetic information by the introduction of a foreign nucleic acid molecule and that the presence or the expression of the foreign nucleic acid molecule leads to a phenotypic change. In this context, phenotypic change preferably means a measurable change of one or more functions of the cells. For example, the genetically modified plant cells of the invention exhibit an increase in the activity of a plastidial ADP/ATP translocator due to the presence or upon expression of the foreign nucleic acid molecule introduced.

In the context of the present invention the term "increase in the activity" means an increase of the expression of a plastidial ADP/ATP translocator gene, an increase in the amount of plastidial ADP/ATP translocator protein and/or an increase in the activity of a plastidial ADP/ATP translocator in the cells.

The increase in the expression can, for example, be determined by measurement of the amount of transcripts encoding ADP/ATP translocator, for example by means of Northern blot analysis. In this context, an increase preferably means an increase in the amount of transcripts in comparison with corresponding non-genetically modified cells by at least 10%, preferably by at least 20%, particularly by at least 50% and particularly preferred by at least 75%. The increase in the amount of ADP/ATP translocator protein can, for example, be determined by Western blot analysis. In this context, an increase preferably means an increase in the amount of ADP/ATP translocator protein in comparison with corresponding non-genetically modified cells by at least 10%, preferably by at least 20%, particularly by at least 50% and particularly preferred by at least 75%.

The activity of the plastidial ADP/ATP translocator can be determined, for example, by isolating the plastids from the corresponding tissue and determining the V_{\max} -values of the ATP import by means of the silicone oil filtration method. The

purification of various plastid types is described in e.g. Neuhaus et al. (Biochem. J. 296 (1993), 395-401). The silicone oil filtration method is described e.g. in Quick et al. (Plant Physiol. 109 (1995), 113-121).

- 5 It was surprisingly found that with plants containing said plant cells with increased activity of the plastidial ADP/ATP translocator the yield of content substances and/or biomass is increased in comparison with corresponding non-modified wild type plants. It was, for example, found that the oil content and/or the starch content in plants according to the invention is increased and/or that also the amylose content of
10 these starches is increased in comparison with non-modified wild type plants.

In this context, the term "wild type plant" refers to plants which serve as starting material for the production of the plants described, i.e. plants of which the genetic information – apart from the genetic modification introduced – is identical to the
15 genetic information of a plant of the invention.

The term "increased yield" means that the portion of content substances, preferably starch or oil in the plant cells of the invention is increased by at least 10%, preferably by at least 20%, more preferably by at least 30% and most preferably by at least 40% in comparison with plant cells of non-modified wild type plants.

20 The term "increased starch content" means that the portion of starch in plant cells according to the invention is increased by at least 10%, preferably by at least 20%, more preferably by at least 30% and most preferably by at least 40% in comparison with plant cells of non-modified wild type plants.

The determination of the starch portion is carried out according to the methods
25 described in the appended Examples.

The term "increased amylose content" means that the amylose content of the starch synthesized in the plant cells of the invention is increased by at least 10%, preferably by at least 20%, more preferably by at least 30% and most preferably by at least 40% in comparison with plant cells of non-modified wild type plants.

30 The amylose content is determined by carrying out the methods described in the appended Examples.

As mentioned above, the plastidial ADP/ATP translocator is a transport protein which is localized in the inner membrane of plastids (Heldt et al, FEBS Lett. 5 (1969), 11-
35 14; Pozueta-Romero et al., Proc. Nat. Acad. Sci. USA 88 (1991), 5769-5773; Neuhaus, Plant Physiol. 101 (1993) 573-578; Schünemann et al., Plant Physiol. 103 (1993), 131-137) and which catalyzes the transport of ATP into the plastids and of ADP out of the plastids. Thus, the plastidial ADP/ATP translocator provides the stroma with cytosolic ATP.

Kampfenkel et al. (FEBS Lett. 374 (1995), 351-355) were the first to isolate a cDNA encoding an ADP/ATP translocator (AATP1) from *Arabidopsis thaliana* (Neuhaus et al., Plant J. 11 (1997), 73-82) and which exhibits a great similarity (66.2% similarity) to the ADP/ATP translocator of the Gram-negative bacterium *Rickettsia prowazekii*.

5 The AATP1-cDNA from *A. thaliana* encodes a strongly hydrophobic protein consisting of 589 amino acids which exhibits 12 potential transmembrane helices (Kampfenkel et al., FEBS Lett. 374 (1995), 351-355). Said cDNA could be functionally expressed in bakers' yeast and *E. coli*. After extraction of the protein and reconstitution in proteoliposomes an increase in the ATP transport rate could be
10 determined (Neuhaus et al., Plant J. 11 (1997), 73-82). By means of antibodies against a peptide fragment of the AATP1 from *A. thaliana* it could be shown that the ADP/ATP translocator AATP1 is localized in the inner chloroplast envelope membrane (Neuhaus et al., Plant J. 11 (1997), 73-82).

The function of the plastidial ADP/ATP translocator for the plant metabolism could
15 not be definitely clarified so far. Various functions have been taken into consideration, e.g. that the supply of the stroma with cytosolic ATP could have an influence on the import of proteins into the plastids, on the amino acid biosynthesis, the fatty acid metabolism or the starch metabolism (Flügge and Hinz, Eur. J. Biochem. 160 (1986), 563-570; Tetlow et al., Planta 194 (1994), 454-460; Hill and
20 Smith, Planta 185 (1991), 91-96; Kleppinger-Sparace et al., Plant Physiol. 98 (1992), 723-727).

The fact that an increase in the activity of the plastidial ADP/ATP translocator leads to an increase in the starch content in the corresponding transgenic plants was, however, completely surprising. Just as surprising was the finding that the increase in
25 the activity of the plastidial ADP/ATP translocator has an effect on the molecular composition of the produced starch. The starch from tubers of potato plants according to the invention, for example, exhibits an increased amylose content in comparison with starches from tubers of non-transformed potato plants.

So far, it has been assumed that the molecular properties of starch are exclusively
30 determined by the interaction of starch-synthesizing enzymes, such as the branching enzymes (E.C. 2.4.1.18), the starch synthases (E.C. 2.4.1.21) and the ADP-glucosepyrophosphorylase (E.C. 2.7.7.27). The fact that the expression of a plastidial transport protein has an influence on the structure of the starch is, however, completely surprising.

35 The plant cells of the invention can be derived from any plant species, i.e. both from monocotyledonous and from dicotyledonous plants. Preferably the plant cells are from agricultural crop plants, i.e. from plants cultivated by humans for the purpose of nutrition or for technical, particularly industrial purposes. Generally preferred are

plant cells from oil- and/or starch-synthesizing or oil- and/or starch-storing plants. Thus, the invention preferably relates to plant cells from starch-synthesizing or starch-storing plants such as cereals (rye, barley, oat, wheat, millet, sago etc.), rice, peas, maize, medullar pea, cassava, potato, rape, soy bean, hemp, flax, sunflower or
5 vegetables (tomato, chicory, cucumber, salad etc.). Plant cells from potato, sunflower, soy bean, rice are preferred. Particularly preferred are plant cells from maize, wheat, rape and rice.

Furthermore, subject-matter of the invention are transgenic plants containing the
10 transgenic plant cells described above. Said plants can be produced, for example, by regeneration from plant cells of the invention. The transgenic plants can, in principle, be plants of any species, i.e. both monocotyledonous and dicotyledonous plants. Preferably, they are useful plants i.e. plants cultivated by humans for the purpose of nutrition or for technical, particularly industrial purposes. These plants can be oil-
15 and/or starch-synthesizing or oil- and/or starch-storing plants. The invention preferably relates to plants such as cereals (rye, barley, oat, wheat, millet, sago etc.), rice, peas, maize, medullar pea, cassava, potato, rape, soy bean, hemp, flax, sunflower or vegetables (tomato, chicory, cucumber, salad etc.). Preferred are potato, sunflower, soy bean, rice. Particularly preferred are maize, wheat, rape and
20 rice.

As mentioned before, it was surprisingly found that in starch-storing plants containing plant cells of the invention with increased activity of the plastidial ADP/ATP translocator the starch content is increased in comparison with wild type plants
25 and/or that also the amylose content of these starches is increased in comparison with corresponding non-modified wild type plants.

Thus, in a preferred embodiment the present invention also relates to starch-storing plants which contain the plant cells of the invention and which exhibit an increased
30 starch content in comparison with non-modified wild type plants and/or an increased amylose content of said starch in comparison with corresponding non-modified wild type plants.

The term "starch-storing plants" comprises all plants with starch-storing tissues such as maize, wheat, rice, potato, rye, barley, oat. Rice, barley and potato are preferred.
35 Particularly preferred are maize and wheat.

In this context, an increase in the "yield" ("increased yield"), an increase in the starch content ("increased starch content"), an increase in the amylose content ("increased amylose content") and the term "wild type plant" are used within the meaning of the

definitions above and are used within the same meaning for the following embodiments of the invention, too. The term "increased yield" preferably means an increase in the production of content substances and/or biomass, in particular, if it is measured by means of fresh weight per plant.

5

Said increase in the yield preferably relates to parts of plants which can be harvested such as seeds, fruits, storage roots, roots, tubers, blossoms, buds, shoots, stems or wood.

10

According to the invention the increase in the yield is at least 3% with regard to the biomass and/or content substances in comparison with corresponding non-transformed plants of the same genotype, if said plants are cultivated under the same conditions, preferably at least 10%, more preferably at least 20% and most preferably at least 30% or even 40% in comparison with wild type plants.

15

Said plants according to the invention have, for example, in comparison with other plants synthesizing starch with increased amylose content such as the *amylose-extender* and the *dull* mutants from maize, the advantage that apart from an increased amylose content they exhibit no reduced but even an increased starch content.

20

Moreover, subject-matter of the present invention are oil-storing plants which contain the plant cells of the invention and which exhibit an increased oil content in comparison with non-modified wild type plant cells, preferably in cells of oil-storing tissue.

25

The term "oil-storing plants" comprises all plants able to store oil such as rape, canola, soy bean, sunflower, maize, peanut, wheat, cotton, oil palms, olive trees and avocado. Preferred are maize, wheat and soy bean. Particularly preferred are rape and canola.

30

The term "increased oil content" means that the oil content in plant cells of the invention is increased by at least 10%, preferably by at least 20%, more preferably by at least 30% and most preferably by at least 40% in comparison with plant cells from non-modified wild type plants.

35

Methods for the determination of the oil content are known to the person skilled in the art and described, for example, by Matthaeus and Bruehl, GIT Labor-Fachz. 43 (1999), 151-152, 154-155; Matthaeus, Laborpraxis 22 (1998), 52-55. The determination of the oil content may also be carried out by non-invasive near IR-spectroscopy which is an analysing method (commonly used in breeding) and was described e.g. by Schulz et al., J. Near Infrared Spectrosc. 6 (1998), A125-A130; Starr et al., J. Agric. Sci. 104 (1985), 317-323.

Plants exhibiting an increased concentration of oil are of great commercial interest. Maize plants, for example, whose grains exhibit a high level of starch but also an increased content of the side product oil are of great interest for the wet milling industry since the side product is of high value. The feed-stuff industry is also
 5 interested in feeding plants with increased oil content since such plants have an increased nutritious value. For the oil plants-processing industry an increase of the oil content means an increase of the efficiency of the oil extracting process.

The present invention further relates to a method for the production of transgenic
 10 plants which, compared to wild type plants, exhibit an increased yield, wherein

- (a) a plant cell is genetically modified by means of introduction of a foreign nucleic acid molecule and the genetic modification leads to an increase of the activity of a plastidial ADP/ATP translocator; and
- (b) a plant is regenerated from the cell; and optionally
- 15 (c) further plants are produced from the plant according to (b).

The present invention further relates to a method for the production of transgenic plants which, compared to wild type plants, exhibit an increased starch content and/or whose starch exhibits an increased amylose content in comparison with
 20 corresponding wild type plants, wherein

- (a) a plant cell is genetically modified by means of introduction of a foreign nucleic acid molecule and the genetic modification leads to an increase of the activity of a plastidial ADP/ATP translocator; and
- (b) a plant is regenerated from the cell; and optionally
- 25 (c) further plants are produced from the plant according to (b).

Moreover, subject-matter of the present invention is a method for the production of transgenic plants which, compared to wild type plants, exhibit an increased oil content, wherein

- 30 (a) a plant cell is genetically modified by means of introduction of a foreign nucleic acid molecule and the genetic modification leads to an increase of the activity of a plastidial ADP/ATP translocator; and
- (b) a plant is regenerated from the cell; and optionally
- (c) further plants are produced from the plant according to (b).

35 For the modification introduced into the plant cell according to step (a) the same applies as has been discussed above regarding the plant cells and plants of the invention.

The regeneration of plants according to step (b) can be carried out according to

methods known to the person skilled in the art.

The generation of further plants according to step (c) of the methods of the invention can be achieved e.g. by vegetative propagation (for example via cuttings, tubers or via callus culture and regeneration of whole plants) or by sexual reproduction.

- 5 Preferably, sexual reproduction takes place in a controlled manner, i.e. selected plants with specific properties are crossed with each other and propagated.

The present invention also relates to the plants obtainable by the method of the invention.

10

The present invention also relates to propagation material of plants according to the invention as well as of the transgenic plants produced according to the methods of the invention which contains genetically modified cells of the invention. In this context, the term propagation material comprises those components of the plant which are suitable for the generation of descendants by means of a vegetative or sexual way. Suitable for vegetative propagation are, for example, cuttings, callus cultures, rhizomes or tubers. Other propagation material comprises, for example, fruit, seeds, seedlings, protoplasts, cell cultures, etc. Preferably, propagation material are tubers, particularly preferred seeds.

20

The present invention further relates to the use of nucleic acid molecules encoding a plastidial ADP/ATP translocator for the production of transgenic plants with an increased yield in comparison with wild type plants.

25

The present invention further relates to the use of nucleic acid molecules encoding a plastidial ADP/ATP translocator for the production of plants which, in comparison with wild type plants, have an increased starch content in the starch-synthesizing and/or -storing tissue, or for the production of plants synthesizing a starch which, compared to starch from wild type plants, exhibits an increased amylose content. The nucleic acid molecules mentioned above in connection with the cells of the invention are preferably used.

30

The present invention further relates to the use of nucleic acid molecules encoding a plastidial ADP/ATP translocator for the production of transgenic plants which, in comparison with wild type plants, have an increased oil content.

35

The present invention further relates to transgenic plant cells which are genetically modified, wherein the genetic modification leads to the decrease of the activity of a plastidial ADP/ATP translocator present endogenously in the plant cell, compared to

non-genetically modified plant cells of corresponding wild type plants.

5 The term "transgenic", as used herein, means that the plant cells of the invention deviate in their genetic information from corresponding non-modified plant cells due to a genetic modification, particularly the introduction of a foreign nucleic acid molecule.

10 In this context, the term "genetically modified" means that the plant cell is modified in its genetic information due to the introduction of a foreign nucleic acid molecule and that the presence or the expression of the foreign nucleic acid molecule leads to a phenotypic change. Phenotypic change preferably means a measurable change of one or more functions of the cell. For example, genetically modified plant cells of the invention exhibit a decrease of the activity of a plastidial ADP/ATP translocator.

15 The production of said plant cells of the invention with a decreased activity of an ADP/ATP translocator can be achieved by various methods known to the person skilled in the art, e.g. by methods leading to an inhibition of the expression of endogenous genes encoding a plastidial ADP/ATP translocator. Such methods include, for example, the expression of a corresponding antisense-RNA, the
20 expression of a sense-RNA for achieving a cosuppression effect, the expression of a correspondingly constructed ribozyme which specifically cleaves transcripts encoding an ADP/ATP translocator or the so-called "in-vivo mutagenesis".

For the reduction of the activity of an ADP/ATP translocator in the cells of the invention preferably an antisense-RNA is expressed.

25 For the expression either a DNA molecule can be used comprising the whole sequence encoding a ADP/ATP translocator including flanking sequences that are possibly present or DNA molecules comprising only parts of the coding sequence, wherein these parts have to be long enough to lead to an antisense-effect in the cells. In general, sequences can be used up to a minimum length of 15 bp, preferably
30 a length of 100-500 bp, for an efficient antisense-inhibition, particularly, sequences with a length of more than 500 bp. DNA molecules shorter than 5000 bp are commonly used, preferably sequences shorter than 2500 bp.

It is also possible to use DNA sequences which have a high degree of homology to the sequences which occur endogenously in the plant cell and which encode a
35 plastidial ADP/ATP translocator. The minimal homology should be higher than approximately 65%. The use of sequences with homologies between 95 and 100% is to be preferred.

Alternatively, the reduction of the ADP/ATP translocator activity in the plant cells of

the invention can also be accomplished by means of a co-suppression effect. The method is known to the person skilled in the art and is described, for example, in Jorgensen (Trends Biotechnol. 8 (1990), 340-344), Niebel et al. (Curr. Top. Microbiol. Immunol. 197 (1995), 91-103), Flavell et al. (Curr. Top. Microbiol. Immunol. 197 (1995), 43-46), Palaqui and Vaucheret (Plant. Mol. Biol. 29 (1995), 149-159), Vaucheret et al. (Mol. Gen. Genet. 248 (1995), 311-317), de Borne et al. (Mol. Gen. Genet. 243 (1994), 613-621) and other sources.

The expression of ribozymes for the reduction of the activity of specific proteins in cells is also known to the person skilled in the art and is described, for example, in EP-B1 0 321 201. The expression of ribozymes in plant cells was described, for example, in Feyter et al. (Mol. Gen. Genet. 250 (1996), 329-338).

Moreover, the reduction of the ADP/ATP translocator activity in the plant cells of the invention can also be achieved by means of the so-called "in vivo mutagenesis", wherein a hybrid RNA-DNA-oligonucleotide ("chimeroplast") is introduced into cells by means of transformation of cells (Kipp, P.B. et al., Poster Session at the "5th International Congress of Plant Molecular Biology, 21.-27. September 1997, Singapore; R. A. Dixon and C. J. Arntzen, Meeting report on "Metabolic Engineering in Transgenic Plants", Keystone Symposia, Copper Mountain, CO, USA, TIBTECH 15 (1997), 441-447; international patent application WO 95/15972; Kren et al., Hepatology 25 (1997), 1462-1468; Cole-Strauss et al., Science 273 (1996), 1386-1389).

A part of the DNA component of the RNA-DNA-oligonucleotide is homologous to a nucleic acid sequence of an endogenous ADP/ATP translocator but, compared to the nucleic acid sequence of the endogenous ADP/ATP translocator, exhibits a mutation or contains a heterologous region which is enclosed by the homologous regions.

By means of base pairing of the homologous regions of the RNA-DNA-oligonucleotide and the endogenous nucleic acid molecule followed by homologous recombination, the mutation or heterologous region contained in the DNA component of the RNA-DNA-oligonucleotide can be transferred into the genome of a plant cell. This leads to a decrease of the activity of the plastidial ADP/ATP translocator.

Thus, subject-matter of the present invention particularly are transgenic plant cells,

- (a) containing a DNA molecule which can lead to the synthesis of an antisense RNA causing a decrease of the expression of endogenous genes which encode a plastidial ADP/ATP translocator; and/or
- (b) containing a DNA molecule which can lead to the synthesis of a co-suppression-RNA causing a decrease of the expression of endogenous genes

- which encode a plastidial ADP/ATP translocator; and/or
- (c) containing a DNA molecule which can lead to the synthesis of a ribozyme which can specifically cleave transcripts of endogenous genes encoding an ADP/ATP translocator; and/or
- 5 (d) which, due to an in vivo mutagenesis, exhibit a mutation or an insertion of a heterologous DNA sequence in at least one endogenous gene encoding a plastidial ADP/ATP translocator, wherein the mutation or insertion causes a decrease of the expression of the gene or the synthesis of an inactive transporter molecule.

10

The term "decrease of the activity" in the present invention means a decrease of the expression of endogenous genes encoding an ADP/ATP translocator, a reduction of the amount of ADP/ATP translocator protein in the cells and/or a decrease of the biological activity of the ADP/ATP translocator protein in the cells.

- 15 The decrease of the expression can be determined, for example, by measuring the amount of transcripts encoding the ADP/ATP translocator, e.g. by Northern blot analysis. A decrease preferably means a decrease of the amount of transcripts in comparison with genetically non-modified cells by at least 30%, preferably by at least 50%, more preferably by at least 70%, particularly preferred by at least 85% and
- 20 most preferably by at least 95%.

- The decrease of the amount of ADP/ATP translocator protein can be determined, for example, by means of Western blot analysis. A decrease preferably means a decrease of the amount of ADP/ATP translocator protein in comparison with corresponding genetically non-modified cells by at least 30%, preferably by at least
- 25 50%, more preferably by at least 70%, particularly preferred by at least 85% and most preferably by at least 95%.

- Surprisingly, it was found that the starch content of plant cells which have a decreased expression and thus a decreased activity of the plastidial ADP/ATP
- 30 translocator, compared to corresponding non-modified plant cells from wild type plants, is reduced and that also the amylose content of these starches, compared to corresponding non-modified plant cells from wild type plants, is reduced. The fact that the starches of the plants of the invention have a modified structure is particularly surprising since it has been assumed so far that the molecular properties
- 35 of starches are exclusively determined by the interaction of starch-synthesizing enzymes such as the branching enzymes (E.C. 2.4.1.18) and the starch synthases (E.C. 2.4.1.21). It is completely surprising that the expression of a plastidial transport protein influences the structure of starch.

The term "decreased starch content" in the present invention means that the content

of starch in plant cells of the invention is reduced by at least 15%, preferably by at least 30%, more preferably by at least 40% and most preferably by at least 50% in comparison with plant cells of non-modified wild type plants. The starch content is determined according to the methods described in the Examples.

- 5 The term "decreased amylose content" means that the content of amylose in the plant cells of the invention, in comparison with plant cells of non-modified wild type plants, is reduced by at least 10%, preferably by at least 20%, more preferably by at least 30% and most preferably by at least 40%. The amylose content is determined according to the methods described in the Examples.
- 10 The term "wild type plant" has the above-defined meaning.

The plant cells of the invention can derive from any plant species, i.e. both from monocotyledonous and dicotyledonous plants. Preferably these are plant cells from agricultural crop plants, i.e. from plants cultivated by humans for the purpose of
15 nutrition or for technical, particularly industrial purposes. Preferably, thus, the invention relates to plant cells from starch-synthesizing or starch-storing plants such as cereals (rye, barley, oat, wheat, millet, sago etc.), rice, pea, maize, medullar pea, cassava, potato, tomato, rape, soy bean, hemp, flax, sunflower, cow pea and arrowroot. Particularly preferred are plant cells from potato.

- 20 Moreover, subject-matter of the invention are transgenic plants containing the transgenic plant cells described above. Said plants can be produced by regeneration from the plant cells of the invention. The transgenic plants can, in principle, be plants of any plant species, i.e. both monocotyledonous and dicotyledonous plants.
- 25 Preferably they are plant cells from agricultural crop plants, i.e. from plants cultivated by humans for the purpose of nutrition or for technical, particularly industrial purposes. Preferably these are starch-synthesizing or starch-storing plants such as cereals (rye, barley, oat, wheat, millet, sago etc.), rice, pea, maize, medullar pea, cassava, potato, tomato, rape, soy bean, hemp, flax, sunflower, cow pea and
30 arrowroot. Particularly preferred is potato.

Said plants of the invention synthesize a starch which, compared to starch from corresponding wild type plants, exhibits a reduced amylose content. The terms
35 "reduction of the amylose content" and "wild type plants" are defined as described above.

Furthermore, the present invention also relates to a method for the production of transgenic plants whose starch, compared to starch from corresponding wild type plants, exhibits a reduced amylose content wherein

- (a) a plant cell is genetically modified by means of introduction of a foreign nucleic acid molecule and the genetic modification leads to a decrease of the activity of a plastidial ADP/ATP translocator present endogenously in plant cells; and
- (b) a plant is regenerated from the cell produced according to step (a); and
- 5 optionally
- (c) further plants are produced from the plant produced according to step (b).

10 For the modification introduced into the plant cell according to step (a) the same applies as was discussed earlier in connection with the plant cells and plants of the invention.

The regeneration of plants according to step (c) can be carried out according to methods known to the person skilled in the art.

15 The production of further plants according to step (c) of the method of the invention can, for example, be achieved by vegetative propagation (for example via cuttings, tubers or via callus culture and regeneration of whole plants) or by sexual propagation. Preferably, sexual propagation takes place in a controlled manner, i.e. selected plants with specific properties are crossed with each other and propagated.

20 In a preferred embodiment the method of the invention is used for the production of transgenic potato plants.

The present invention also relates to plants obtainable by the method of the invention.

25 The present invention also relates to propagation material of plants according to the invention as well as of the transgenic plants produced according to the methods of the invention which contains genetically modified cells of the invention. In this context, the term propagation material comprises those components of the plant which are suitable for the generation of descendants by means of a vegetative or

30 sexual way. Suitable for vegetative propagation are, for example, cuttings, callus cultures, rhizomes or tubers. Other propagation material comprises, for example, fruit, seeds, seedlings, protoplasts, cell cultures, etc. Preferably, propagation material are seeds, particularly preferred tubers.

35 Furthermore, the present invention relates to the use of nucleic acid molecules encoding a plastidial ADP/ATP translocator, of complements thereof or of parts of said molecules for the production of plants synthesizing a starch with, in comparison with starch from wild type plants, reduced amylose content. Preferably, the nucleic acid molecules mentioned above in connection with the plant cells of the invention

exhibiting an increased ADP/ATP translocator activity are to be used.

A variety of techniques are at disposal for the introduction of DNA in a plant host cell. These techniques comprise the transformation of plant cells with T-DNA using
 5 *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes* as transformation agent, the fusion of protoplasts, the injection, the electroporation of DNA, the introduction of the DNA via the biolistic approach and other possibilities.

The use of *Agrobacteria*-mediated transformation of plant cells has been analysed in detail and was described sufficiently in EP 120516; Hoekema, In: The Binary Plant
 10 Vector System Offsetdrukkerij Kanters B.V., Alblasterdam (1985), Chapter V; Fraley et al., Crit. Rev. Plant Sci. 4, 1-46 and An et al., EMBO J. 4 (1985), 277-287. For the transformation of potato, see e.g. Rocha-Sosa et al., EMBO J. 8 (1989), 29-33.

The transformation of monocotyledonous plants by means of *Agrobacterium*-based
 15 vectors was also described (Chan et al., Plant Mol. Biol. 22 (1993), 491-506; Hiei et al., Plant J. 6 (1994) 271-282; Deng et al., Science in China 33 (1990), 28-34; Wilmink et al., Plant Cell Reports 11 (1992), 76-80; May et al., Bio/Technology 13 (1995), 486-492; Connor and Domisse, Int. J. Plant Sci. 153 (1992), 550-555; Ritchie et al., Transgenic Res. 2 (1993), 252-265). An alternative system for the transformation of monocotyledonous plants is the transformation via the biolistic
 20 approach (Wan and Lemaux, Plant Physiol. 104 (1994), 37-48; Vasil et al., Bio/Technology 11 (1993), 1553-1558; Ritala et al., Plant Mol. Biol. 24 (1994), 317-325; Spencer et al., Theor. Appl. Genet. 79 (1990), 625-631), the protoplast transformation, the electroporation of partially permeabilized cells, the introduction of DNA by means of glass fibres. The transformation of maize, in particular, is
 25 described in the literature several times (see e.g. WO 95/06128, EP 0513849, EO 0465875, EP 292435; Fromm et al., Biotechnology 8 (1990), 833-844; Gordon-Kamm et al., Plant Cell 2 (1990), 603-618; Koziel et al., Biotechnology 11 (1993), 194-200; Moroc et al., Theor. Appl. Genet. 80 (1990), 721-726).

The successful transformation of other cereals has also been described, e.g. for
 30 barley (Wan and Lemaux, loc. cit., Ritala et al., loc. cit., Krens et al., Nature 296 (1982), 72-74) and for wheat (Nehra et al., Plant J. 5 (1994), 285-297).

For the expression of the nucleic acid molecules encoding an ADP/ATP translocator in sense- or antisense-orientation in plant cells said nucleic acid molecules are
 35 preferably linked with regulatory DNA elements which ensure the transcription in plant cells. Said elements include, in particular, promoters. Generally, any promoter active in plant cells is suitable.

The promoter can be chosen in such a way that the expression takes place constitutively or only in a specific tissue, at a specific point in time of the plant

development or at a point in time determined by external factors. Both with regard to the plant and with regard to the nucleic acid molecule, the promoter can be homologous or heterologous.

Suitable promoters are e.g. the promoter of the 35S RNA of the cauliflower mosaic virus and the ubiquitin promoter from maize for a constitutive expression, the patatin gene promoter B33 (Rocha-Sosa et al., EMBO J. 8 (1989), 23-29) for a tuber-specific expression in potato and a promoter ensuring an expression only in photosynthetically active tissue, e.g. the ST-LS1-promoter (Stockhaus et al., Proc. Natl. Acad. Sci. USA 84 (1987), 7943-7947; Stockhaus et al., EMBO J. 8 (1989), 2445-2451) or, for an endosperm-specific expression, the HMG promoter from wheat, the USP promoter, the phaseolin promoter, promoters from zein genes from maize (Pedersen et al., Cell 29 (1982), 1015-1026; Quatroccio et al., Plant Mol. Biol. 15 (1990), 81-93), glutelin promoter (Leisy et al., Plant Mol. Biol. 14 (1990), 41-50; Zheng et al., Plant J. 4 (1993), 357-366; Yoshihara et al., FEBS Lett. 383 (1996), 213-218) or shrunken-1 promoter (Werr et al., EMBO J. 4 (1985), 1373-1380). Promoters which are activated only at a point in time determined by external factors can, however, also be used (see for example WO 9307279). In this context, promoters of heat-shock proteins allowing for a simple induction can particularly be of interest. Moreover, seed-specific promoters can be used such as the USP promoter from *Vicia faba* which ensures a seed-specific expression in *Vicia faba* and other plants (Fiedler et al., Plant Mol. Biol. 22 (1993), 669-679; Bäumlein et al., Mol. Gen. Genet. 225 (1991), 459-467).

The aforementioned embodiments with the endospecific¹ promoters are suitable, in particular, for increasing the starch content in the endosperm. In contrast thereto, the use of embryo-specific promoters is of interest, in particular, for increasing the oil content since, as a rule, oil is mainly stored in the embryo.

Thus, preferably a promoter is used according to the present invention which ensures the expression in the embryo or in the seed. In a preferred embodiment of the invention the promoter is the globulin-1 (glb1) promoter from maize (Styer and Cantliffe, Plant Physiol. 76 (1984), 196-200). In another embodiment of the invention the embryo-specific promoter is from plants, preferably from *Cuphea lanceolata*, *Brassica rapa* or *Brassica napus*. Particularly preferred are the promoters pCIFatB3 and pCIFatB4 (WO 95/07357). These are promoters of the genes CIFatB3 and CIFatB4, respectively, which have already successfully been used in transgenic rape for the biosynthesis of medium-chain fatty acids and, thus, have a suitable expression window for the solution of the present problem.

¹ Translator's note: should read "endosperm-specific"

In another preferred embodiment the pCIGPDH promoter (WO 95/06733), the napin (described, for example, by Kridl, Seed Sci. Res. 1 (1991), 209-219; Ellerstrom et al., Plant Mol. Biol. 32 (1996), 1019-1027; Stalberg et al., Planta 199 (1996), 515-519) or the oleosin promoter (described, for example, by Keddie, Plant Mol. Biol. 24 (1994), 327-340; Plant et al., Plant Mol. Biol. 25 (1994), 193-205) is used.

Moreover, a termination sequence can be present which serves the correct termination of the transcription and the addition of a poly-A-tail to the transcript regarded as having a function in stabilizing the transcripts. Said elements are described in the literature (see, e.g., Gielen et al., EMBO J. 8 (1989), 23-29) and are interchangeable as desired.

The transgenic plant cells and plants of the invention synthesize, preferably due to the increase or decrease of the activity of a plastidial ADP/ATP translocator, a starch which, compared to synthesized starch in wild type plants, is modified in its physio-chemical properties, in particular the amylose/amylopectin ratio. In particular, said starch can, compared to wild type starch, be modified with regard to the viscosity and/or the gel formation properties of gels of said starch

Thus, the present invention relates to methods for the production of a modified starch comprising the step of extraction of the starch from one of the above-described plants and/or from starch-storing parts of said plant. Preferably, said method comprises also the step of harvesting the cultivated plants and/or starch-storing parts of said plants before the extraction of starch and further, particularly preferred, the step of cultivating the plants of the invention before harvesting. Methods for the extraction of the starch from plants or the starch-storing parts of plants are known to the person skilled in the art. Moreover, methods for the extraction of the starch from various other starch-storing plants are described, e.g. in "Starch: Chemistry and Technology (eds.: Whistler, BeMiller and Paschall (1994), 2nd edition, Academic Press Inc. London Ltd; ISBN 0-12-746270-8; see e.g. chapter XII, page 412-468: maize and sorghum starch: production; by Watson; chapter XIII, page 469-479: starches from tapioca, arrowroot and sago: production; by Corbishley and Miller; chapter XIV, page 491-506: starch from wheat: production, modification and uses; by Knight and Oson; and chapter XVI, page 507 to 528: starch from rice: production and uses; by Rohmer and Klem; starch from maize: Eckhoff et al., Cereal Chem. 73 (1996) 54-57, the extraction of starch from maize to industrial standard is usually achieved by so-called 'wet milling'). Appliances usually used for methods for the extraction of starch from plant material are separators, decanters, hydrocyclones, spray dryers and fluidized bed dryers.

Furthermore, subject-matter of the present invention is starch obtainable from the transgenic plant cells, plants and propagation material of the invention and starch obtainable by the method of the invention described above.

The starches of the invention can be modified according to methods known to the person skilled in the art and are suitable for various uses in the foodstuff or non-foodstuff industry in unmodified or modified form.

In principle, possibilities of use can be divided into two large areas. One area comprises hydrolysis products of the starch, mainly glucose and glucan building blocks obtained via enzymatic or chemical methods. They serve as starting material for further chemical modifications and processes such as fermentation. For a reduction of costs the simplicity and inexpensive carrying out of a hydrolysis method can be of importance. At present, the method is essentially enzymatic with use of amyloglucosidase. It would be possible to save costs by reducing use of enzymes. This could be achieved by changing the structure of the starch, e.g. surface enlargement of the grain, easier digestibility due to low branching degree or a steric structure limiting the accessibility for the enzymes used.

The other area where starch is used as so-called native starch due to its polymeric structure can be subdivided into two further fields of application:

1. Use in foodstuffs

Starch is a classic additive for various foodstuffs, in which it essentially serves the purpose of binding aqueous additives and/or causes an increased viscosity or an increased gel formation. Important characteristic properties are flowing and sorption behaviour, swelling and pastification temperature, viscosity and thickening performance, solubility of the starch, transparency and paste structure, heat, shear and acid resistance, tendency to retrogradation, capability of film formation, resistance to freezing/thawing, digestibility as well as the capability of complex formation with e.g. inorganic or organic ions.

2. Use in non-foodstuffs

The other major field of application is the use of starch as an adjuvant in various production processes or as an additive in technical products. The major fields of application for the use of starch as an adjuvant are, first of all, the paper and cardboard industry. In this field, the starch is mainly used for retention (holding back solids), for sizing filler and fine particles, as solidifying

substance and for dehydration. In addition, the advantageous properties of starch with regard to stiffness, hardness, sound, grip, gloss, smoothness, tear strength as well as the surfaces are utilized.

5 2.1 Paper and cardboard industry

Within the paper production process, a differentiation can be made between four fields of application, namely surface, coating, mass and spraying.

10 The requirements on starch with regard to surface treatment are essentially a high degree of brightness, corresponding viscosity, high viscosity stability, good film formation as well as low formation of dust. When used in coating the solid content, a corresponding viscosity, a high capability to bind as well as a high pigment affinity play an important role. As an additive to the mass rapid, uniform, loss-free dispersion, high mechanical stability and complete retention in the paper pulp are of importance. When using the starch in spraying,
15 corresponding content of solids, high viscosity as well as high capability to bind are also significant.

2.2 Adhesive industry

20 A major field of application is, for instance, in the adhesive industry, where the fields of application are subdivided into four areas: the use as pure starch glue, the use in starch glues prepared with special chemicals, the use of starch as an additive to synthetic resins and polymer dispersions as well as the use of starches as extenders for synthetic adhesives. 90% of all starch-based adhesives are used in the production of corrugated board, paper sacks
25 and bags, composite materials for paper and aluminum, boxes and wetting glue for envelopes, stamps, etc.

2.3 Textiles and textile care products

30 Another possible use as adjuvant and additive is in the production of textiles and textile care products. Within the textile industry, a differentiation can be made between the following four fields of application: the use of starch as a sizing agent, i.e. as an adjuvant for smoothing and strengthening the burring behaviour for the protection against tensile forces active in weaving as well as for the increase of wear resistance during weaving, as an agent for textile

improvement mainly after quality-deteriorating pretreatments, such as bleaching, dying, etc., as thickener in the production of dye pastes for the prevention of dye diffusion and as an additive for warping agents for sewing yarns.

5

2.4 Building industry

10

Furthermore, starch may be used as an additive in building materials. One example is the production of gypsum plaster boards, in which the starch mixed in the thin plaster pastifies with the water, diffuses at the surface of the gypsum board and thus binds the cardboard to the board. Other fields of application are admixing it to plaster and mineral fibers. In ready-mixed concrete, starch may be used for the deceleration of the sizing process.

2.5 Ground stabilisation

15

Furthermore, the starch is advantageous for the production of means for ground stabilisation used for the temporary protection of ground particles against water in artificial earth shifting. According to state-of-the-art knowledge, combination products consisting of starch and polymer emulsions can be considered to have the same erosion- and encrustation-reducing effect as the products used so far; however, they are considerably less expensive.

20

2.6 Use in plant protectives and fertilizers

25

Another field of application is the use of starch in plant protectives for the modification of the specific properties of these preparations. For instance, starch is used for improving the wetting of plant protectives and fertilizers, for the dosed release of the active ingredients, for the conversion of liquid, volatile and/or odorous active ingredients into microcrystalline, stable, deformable substances, for mixing incompatible compositions and for the prolongation of the duration of the effect due to a reduced disintegration.

30

2.7 Drugs, medicine and cosmetics industry

Starch may also be used in the fields of drugs, medicine and in the cosmetics industry. In the pharmaceutical industry, starch may be used as a binder for tablets or for the dilution of the binder in capsules. Furthermore, starch is

suitable as disintegrant for tablets since, upon swallowing, it absorbs fluid and after a short time it swells so much that the active ingredient is released. For qualitative reasons, medical lubricating and vulnerary dusting powders are further fields of application. In the field of cosmetics, the starch may for example be used as a carrier of powder additives, such as scents and salicylic acid. A relatively extensive field of application for the starch is toothpaste.

2.8 Starch as an additive in coal and briquettes

Starch can also be used as an additive in coal and briquettes. By adding starch, coal can be quantitatively agglomerated and/or briquetted in high quality, thus preventing premature disintegration of the briquettes. Barbecue coal contains between 4 and 6% added starch, calorated coal between 0.1 and 0.5%. Furthermore, starch is suitable as a binding agent since adding it to coal and briquette can considerably reduce the emission of toxic substances.

2.9 Processing of ore and coal slurry

Furthermore, starch may be used as a flocculant in the processing of ore and coal slurry.

2.10 Additive for casting materials

Another field of application is the use as an additive to process materials in casting. For various casting processes cores produced from sands mixed with binding agents are needed. Nowadays, the most commonly used binding agent is bentonite mixed with modified starches, mostly swelling starches.

The purpose of adding starch is increased flow resistance as well as improved binding strength. Moreover, swelling starches may fulfil more prerequisites for the production process, such as dispersability in cold water, rehydratisability, good mixability in sand and high capability of binding water.

2.11 Rubber industry

In the rubber industry starch may be used for improving the technical and optical quality. Reasons for this are improved surface gloss, grip and appearance. For this purpose, starch is dispersed on the sticky rubberised

surfaces of rubber substances before the cold vulcanization. It may also be used for improving the printability of rubber.

2.12 Production of leather substitutes

- 5 Another field of application for modified starch is the production of leather substitutes.

2.13 Starch in synthetic polymers

- 10 In the plastics market the following fields of application are emerging: the integration of products derived from starch into the processing process (starch is only a filler, there is no direct bond between synthetic polymer and starch) or, alternatively, the integration of products derived from starch into the production of polymers (starch and polymer form a stable bond).

15

The use of the starch as a pure filler cannot compete with other substances such as talcum. This situation is different when the specific starch properties become effective and the property profile of the end products is thus clearly changed. One example is the use of starch products in the processing of thermoplastic materials, such as polyethylene. Thereby, starch and the synthetic polymer are combined in a ratio of 1 : 1 by means of coexpression to form a 'master batch', from which various products are produced by means of common techniques using granulated polyethylene. The integration of starch in polyethylene films may cause an increased substance permeability in hollow bodies, improved water vapor permeability, improved antistatic behaviour, improved anti-block behaviour as well as improved printability with aqueous dyes.

20

25

30

Another possibility is the use of the starch in polyurethane foams. Due to the adaptation of starch derivatives as well as due to the optimisation of processing techniques, it is possible to specifically control the reaction between synthetic polymers and the hydroxy groups of the starch. The results are polyurethane films having the following property profiles due to the use of starch: a reduced coefficient of thermal expansion, decreased shrinking behaviour, improved pressure/tension behaviour, increased water vapour permeability without a change in water acceptance, reduced flammability and cracking density, no drop off of combustible

parts, no halides and reduced aging. Disadvantages that presently still exist are reduced pressure and impact strength.

Product development of film is not the only option. Also solid plastics products, such as pots, plates and bowls can be produced by means of a starch content of more than 50%. Furthermore, the starch/polymer mixtures offer the advantage that they are much easier biodegradable.

Furthermore, due to their extreme capability to bind water, starch graft polymers have gained utmost importance. These are products having a backbone of starch and a side lattice of a synthetic monomer grafted on according to the principle of radical chain mechanism. The starch graft polymers available nowadays are characterised by an improved binding and retaining capability of up to 1000 g water per g starch at a high viscosity. These super absorbers are used mainly in the hygiene field, e.g. in products such as nappies and sheets, as well as in the agricultural sector, e.g. in seed pellets.

What is decisive for the use of the novel starch modified by recombinant DNA techniques are, on the one hand, structure, water content, protein content, lipid content, fiber content, ashes/phosphate content, amylose/amylopectin ratio, distribution of the relative molar mass, branching degree, granule size and shape as well as crystallization, and on the other hand, the properties resulting in the following features: flow and sorption behaviour, pastification temperature, viscosity, thickening performance, solubility, paste structure, transparency, heat, shear and acid resistance, tendency to retrogradation, capability of gel formation, resistance to freezing/thawing, capability of complex formation, iodine binding, film formation, adhesive strength, enzyme stability, digestibility and reactivity.

The production of modified starch by genetically operating with a transgenic plant may modify the properties of the starch obtained from the plant in such a way as to render further modifications by means of chemical or physical methods superfluous. On the other hand, the starches modified by means of recombinant DNA techniques might be subjected to further chemical modification, which will result in further improvement of the quality for certain of the above-described fields of application. These chemical modifications are principally known. These are particularly modifications by means of

- heat treatment
- acid treatment
- oxidation and
- esterification

5

leading to the formation of phosphate, nitrate, sulfate, xanthate, acetate and citrate starches. Other organic acids may also be used for the esterification:

- formation of starch ethers

10 starch alkyl ether, O-allyl ether, hydroxylalkyl ether, O-carboxymethyl ether, N-containing starch ethers, P-containing starch ethers and S-containing starch ethers.

- formation of branched starches

15

- formation of starch graft polymers.

Figure 1 schematically illustrates the plasmid pJT31 (AATP1 (*Arabidopsis thaliana*) sense);

20

Figure 2 schematically illustrates the plasmid pJT32 (AATP1 (*Solanum tuberosum*) antisense);

Figure 3 shows the comparison of the amino-acid sequence of the AATP2¹ from *Arabidopsis thaliana* with the AATP1 (A. *thaliana*)^(SEQ ID NO: 5) and a homologous protein from *Rickettsia prowazekii*_v (Williamson et al., Gene 80 (1989), 269-278);
(SEQ ID NO: 6)
(SEQ ID NO: 7)

25

Figure 4 hydropathy analysis of AATP2 (A. *thaliana*), AATP1 (A. *thaliana*) and the *Rickettsia* ADP/ATP translocator carried out according to the method of von Heijne et al. (Eur. J. Biochem. 180 (1989), 535-545)

30

Figure 5 shows a Northern blot analysis of the expression of the AATP1 (*Solanum tuberosum*) in leaf and tuber of ADP/ATP translocator antisense plants.

35

Figure 6 shows a Northern blot analysis of the expression of the AATP1

(*Arabidopsis thaliana*) in leaf and tuber of ADP/ATP translocator overexpression plants.

Figure 7 Schematic map of the cassette pTE200 for the embryo-specific gene expression. EcoRI, SmaI, BamHI, XhoI, NotI, XbaI, SacI, KpnI, ApaI, Sall and SfiI mark recognition sites for restriction endonucleases. For practical reasons, SfiI (A) and SfiI (B) differ in the variable nucleotide sequence within the recognition sequence. The abbreviations encode as follows: PCIFatB4 = CIFatB4 promoter, tCIFatB4 = CIFatB4 terminator, amp = bacterial resistance against ampicillin, ColE1 ori = "origin of replication" from the plasmid ColE1, f1(-) ori = "origin of replication" from the phage f1.

Figure 8 Schematic map of the ADP/ATP translocator expression cassette pTE208: this derivative of the vector pTE200 (Figure 7) carries a cDNA coding for a plastidial ADP/ATP translocator from *Solanum tuberosum* in sense-orientation.

Figure 9 Schematic map of the binary vector pMH000-0. SfiI, Sall, ClaI, HindIII, EcoRI, NsiI, SmaI, BamHI, SpeI, NotI, KpnI, BglII, ApaI, XhoI, XbaI and BstEII mark recognition sites for restriction endonucleases. SfiI (A) and SfiI (B) differ in the variable nucleotide sequence of their recognition sequence as stated. This is the reason why a recircularization of the starting plasmid is prevented after SfiI cleavage and a directed insertion of the expression cassette from the pTE200 derivative is possible. The abbreviations encode as follows: RB, LB = right and left border region, t35S – termination signal of the 35S *rna* gene from CaMV, pat = phosphotricin acetyl transferase gene, p35S = promoter of the 35S *rna* gene from CaMV, p35S(min) = minimal promoter of the 35S *rna* gene from CaMV, tp-*sul* = sulfonamid resistance gene with transit peptide, *tnos* = termination signal of the nopaline synthase gene, Sm/Sp = bacterial resistance against streptomycin and spectinomycin, parA, parB and parR = plasmid multiplication functions from the plasmid pVS1 with large host area i.a. for *Agrobacterium tumefaciens* and *Escherichia coli*.

The following examples illustrate the invention.

Example 1

Construction of the bacterial expression vector pJT118 and transformation of *E. coli*

5

The AATP2 protein (gene library X94626) from *Arabidopsis thaliana* was N-terminally fused with a "histidine-tag" comprising 10 aminos acids.

For this, the cDNA encoding the whole AATP2 protein from *Arabidopsis thaliana* was isolated by means of a PCR approach. The following oligonucleotide served as
10 sense-primer which, in addition, had a XhoI-restriction site:
cgtgagagatagagagctcgagggtctgattcaaacc (SEQ ID NO: 1); comprising the base pairs 66-102).

An oligonucleotide carrying an additional BamHI-restriction site served as antisense-primer: gatacaacaggaatcctggatgaagc (SEQ ID NO: 2); comprising the base pairs
15 1863-1835). The obtained PCR product was purified by means of an agarose gel, cut with the restriction enzymes XhoI/BamHI and introduced "in frame" in the plasmid pET16b (Novagene, Heidelberg, Germany). This led to the exhibition of a histidine-tag of 10 amino acids at the N-terminus of the cDNA encoding the whole AATP2 protein from *Arabidopsis thaliana* (His-AATP2). Said vector was called pJT118. The
20 sequence of the PCR product was determined by sequencing of both nucleotide strands (Eurogentec). The transformation of *E. coli* C43 (Miroux and Walker, J. Mol. Biol. 260 (1996), 289-298) was carried out according to standard methods.

The *E. coli* strain C43 allows for the heterologous expression of animal (Miroux and Walker, loc. cit.) and plant (Tjaden et al., J. Biol. Chem. (1998) (in press)) membrane
25 proteins.

After transformation of said strain with the vector pJT118 uptake studies with radioactively marked ADP and ATP were carried out. It could be demonstrated by these studies that His-AATP2 can be functionally expressed in *E. coli* C43 in the cytoplasmic membrane of *E. coli*. This showed that AATP2 in fact encodes an
30 ADP/ATP translocator. The presence of a N-terminal histidine-tag leads to an increase (2x-3x) of the transport activity of AATP2 from *A. thaliana* in *E. coli* in comparison with AATP2 without N-terminal his-tag.

35

Example 2

Construction of the plasmid pJT31 and introduction of the plasmid into the genome of potato plants

5

For the construction of a plant transformation vector an EcoRV/BamHI fragment of the AATP1-cDNA from *A. thaliana* (Kampfenkel et al., FEBS Letters 374 (1995), 351-355) with a length of 2230 bp was ligated into the vector pBinAR cut with SmaI/EcoRV and BamHI (Höfgen and Willmitzer, Plant Sci. 66 (1990), 221-230). By

10

insertion of the cDNA fragment an expression cassette is formed (pJT31) which is constructed of the fragments A, B and C as follows (see Figure 1):

Fragment A (540 bp) contains the 35S promoter from the cauliflower mosaic virus.

Fragment B contains, in addition to the flanking regions, the protein-encoding region of an ADP/ATP translocator from *A. thaliana* (AATP1). Said region was isolated as

15

described above and fused in sense-orientation to the 35S promoter in pBinAR. Fragment C (215 bp) contains the polyadenylation signal of the octopine synthase gene from *Agrobacterium tumefaciens*.

The size of the plasmid pJT31 is approximately 14.2 kb.

The plasmid was transferred into potato plants by means of *Agrobacteria* as by

20

Rocha-Sosa et al. (EMBO J. 8 (1989), 23-29). As a result of the transformation transgenic potato plants exhibited an increase of the mRNA of a plastidial ADP/ATP translocator. This was detected by Northern blot analysis (see Fig. 6). RNA was isolated according to standard protocols from leaf and tuber tissue from potato plants. 50µg RNA were separated on an agarose gel (1.5% agarose, 1x MEN puffer, 16.6% formaldehyde). After electrophoresis the RNA was transferred with 20x SSC

25

onto a nylon membrane Hybond N (Amersham, UK) by means of capillary blot. The RNA was fixed on the membrane by means of UV irradiation. The membrane was pre-hybridized for 2 hours in phosphate hybridization buffer (Sambrook et al., loc. cit.) and subsequently hybridized for 10 hours by means of addition of the radioactively

30

labeled probe.

35

Example 3

Construction of the plasmid pJT32 and introduction of the plasmid into the genome of potato plants

5

For the construction of a plant transformation vector a BamHI/NdeI fragment of the coding region of the AATP1-cDNA from *S. tuberosum* (Genbank Y10821) with a length of 1265 bp was ligated into the vector pBinAR (Höfgen and Willmitzer, Plant Sci. 66 (1990), 221-230) cut with SmaI/NdeI and BamHI.

10 By insertion of the cDNA fragment an expression cassette is formed which is constructed of the fragments A, B and C as follows (see Fig. 2):

Fragment A (540 bp) contains the 35S promoter from the cauliflower mosaic virus.

Fragment B contains a region of an ADP/ATP translocator from *S. tuberosum* (AATP1 S.t.) with a length of 1265 bp. This region was fused in antisense-

15 orientation to the 35S promoter in pBinAR.

Fragment C (215 bp) contains the polyadenylation signal of the octopine synthase gene from *Agrobacterium tumefaciens*.

The size of the plasmid pJT32 is approximately 13.3 kb.

20 The plasmid was transferred into potato plants by means of *Agrobacteria* as by Rocha-Sosa et al. (EMBO J. 8 (1989), 23-29).

As a result of the transformation transgenic potato plants exhibited a decrease of the mRNA of a plastidial ADP/ATP translocator. This was detected by Northern blot analysis (see Fig. 5). RNA was isolated according to standard protocols from leaf and tuber tissue from potato plants. 50µg RNA were separated on an agarose gel (1.5%

25 agarose, 1x MEN puffer, 16.6% formaldehyde). After electrophoresis the RNA was transferred with 20x SSC onto a nylon membrane Hybond N (Amersham, UK) by means of capillary blot. The RNA was fixed on the membrane by means of UV irradiation. The membrane was pre-hybridized for 2 hours in phosphate hybridization buffer (Sambrook et al., loc. cit.) and subsequently hybridized for 10 hours by means
30 of addition of the radioactively labeled probe.

Example 4

Analysis of the starch, amylose and sugar content of transgenic potato plants

- 5 The determination of the content of soluble sugars was carried as described by Lowry and Passonneau in "A Flexible System of Enzymatic Analysis", Academic Press, New York, USA (1972). The determination of the starch content was carried out as described by Batz et al. (Plant Physiol. 100 (1992), 184-190).

10 Table 1:

line/genotype	starch in ($\mu\text{molC6units/g}$ fresh weight)	soluble sugars in ($\mu\text{mol/g}$ fresh weight)
Desiree/WT	1094.0	26.49
654/antisense-AATP1 (<i>S. tuberosum</i>)	574.2	42.52
594/antisense-AATP1 (<i>S. tuberosum</i>)	630.2	48.76
595/antisense-AATP1 (<i>S. tuberosum</i>)	531.4	45.92
676/antisense-AATP1 (<i>S. tuberosum</i>)	883.0	40.60
62/sense-AATP1 (<i>A. thaliana</i>)	1485.0	30.65
98/sense-AATP1 (<i>A. thaliana</i>)	1269.0	18.28
78/sense-AATP1 (<i>A. thaliana</i>)	995.0	20.50

The determination of the amylose content was carried out as described by Hovenkamp-Hermelink et al. (Potato Res. 31 (1988), 241-246):

15

20

line/genotype	% amylose
Desiree/WT	18.8
654/antisense-AATP1 (<i>S. tuberosum</i>)	15.5
594/antisense-AATP1 (<i>S. tuberosum</i>)	14.3
595/antisense-AATP1 (<i>S. tuberosum</i>)	18.0
676/antisense-AATP1 (<i>S. tuberosum</i>)	11.5
62/sense-AATP1 (<i>A. thaliana</i>)	27.0
98/sense-AATP1 (<i>A. thaliana</i>)	22.7
78/sense-AATP1 (<i>A. thaliana</i>)	24.5

Example 5

5

Production of an expression cassette and transformation of rape plants

- The expression cassette pTE200 in a pBluescript derivative (Short et al., Nucl. Acid Res.16, (1988), 7583-7600) carries the promoter and terminator sequences of the
- 10 thioesterase gene CIfatB4 (GenBank accession: AJ131741) from *Cuphea lanceolata* and suitable polylinker sequences for the insertion of various useful genes. Peripheral SfiI recognition sites with non-compatible nucleotides in the variable recognition regions allow for a directed transfer of the whole expression cassette including the useful gene into the corresponding restriction sites of the binary plasmid
- 15 vector pMH000-0, a further development of pLH9000 (Hausmann and Töpfer, (1999): 9th chapter: "Entwicklung von Plasmid-Vektoren" in Bioengineering für Rapssorten nach Maß, D. Brauer, G. Röbbelen and R. Töpfer (eds.), Vorträge für Planzenzüchtung, Volume 45, 155-172), and prevent recircularization of the DNA in the recipient vector.
- 20 For the production of the expression cassette pTE200, first, a Sall-BbvI fragment carrying a promoter with an approximate length of 3.3 kb was isolated from the genomic clone CITEg16 (WO95/07357) carrying the complete CIfatB4 gene from *C. lanceolata*. In order to achieve this, the BbvI restriction site at the 3'-end of the promoter was opened and modified in such a way that the fragment could then be

taken up by the pBluescript (stratagene) cut with Sall and SmaI. An internal EcoRI restriction site of the fragment located 1211 nucleotides 5' was deleted by being opened, modified by means of T4 polymerase and subsequently closed again.

The terminator sequence was amplified by means of the polymerase chain reaction and specific oligonucleotide primers at the CITEg16 matrix (WO95/07357) and provided with various polylinker restriction sites (MCS) via the primers. The sequences of the primers are:

5'GAATTCCTGCAGCCCGGGGGATCCACTAGTCTCGAGAAGTGGCTGGGGGCCT
TTCC3' (SEQ ID NO: 3)= 5'-primer: (MCS: EcoRI, PstI, SmaI, BamHI, SpeI XhoI;
CIFatB4 terminator: from pos. 35 – 56) and

5'TCTAGAGGCCAAGGCGGCCGCTTCAACGGACTGCAGTGC3' (SEQ ID NO: 4)
= 3'-primer: CIFatB4 terminator: from pos. 22 – 39, MCS: NotI, StyI, SfiI, XbaI. The
amplificate was cut with EcoRI and NotI and inserted into the corresponding
restriction sites of the pBlueSfi BA (Hausmann and Töpfer, see above). The fragment
carrying the promoter was opened with BamHI, modified and subsequently cut with
Sall to place it in the pBlueSfi BA vector via Sall and modified HindIII restriction site in
front of the terminator. The result is the expression cassette pTE200 (see Figure 7).

For the construction of a plant transformation vector an EcoRI fragment of the AATP1
cDNA from *Solanum tuberosum* (pTM1, Tjaden et al., The Plant Journal 16 (1998)
531-540) with a length of 2270 bp was ligated into the vector pTE200 opened with
EcoRI. The orientation was controlled by means of restriction digest. The result was
the plasmid pTE208 (Figure 8). In the following step, the SfiI fragment from pTE208
was inserted into the polylinker restriction sites of the binary vector pMH000-0
(Figure 9) in a directed manner. The result was the vector pMH 0208.

The binary plasmid vector pMH000-0 has been developed further from pLH9000
(Hausmann and Töpfer, see above) with alternative selection markers for the plant
transformation. The sulfonamide gene (sul) was isolated together with the signal
peptide sequence (tp) for plastidial import of the small subunit of the
ribulosebiphosphate carboxylase from the precursor plasmid of pS001 (Reiss et al.,
Proc. Natl. Acad. Sci. USA 93, (1996), 3094-3098) after modification of the Asp718-
to the XhoI-restriction site. The XhoI-Sall fragment was inserted into the XhoI- and
BamHI-restriction sites of a pBluescript derivative in front of the terminator of the
nopaline synthase gene (pAnos) after modification of Sall and BamHI. In a subsequent
three fragment ligation the resulting tpsul-pAnos fragment (XhoI – XbaI) and the XhoI
– HindIII fragment from pRT103pat (Töpfer et al., Methods in Enzymol. 217, (1993),
66-78) were united with the plasmid pK18 (Pridmore, Gene 56, (1987), 309-312)
opened by means of HindIII and XbaI. As a result the gene for the phosphinotricin
acetyltransferase with the terminator of the CaMV35S rna gene from pRT103pat was
placed in opposite orientation to the tpsul-pAnos unit. A dual promoter of the

CaMV35S rna gene as XhoI fragment from a descendant of pROA93 (Ott et al., Mol. Gen. Genet. 221, (1990), 121-124) was inserted into the XhoI restriction site between the resistance-mediating gene sequences to complete said double selection unit (for resistance against the herbicide Basta and the sulfonamide sulfadiazin). After
5 corresponding modifications in the adjacent polylinker the resulting dual selection cassette was exchanged by means of XbaI and HindIII against the kanamycin cassette in the pLH9000 precursor plasmid (Hausmann and Töpfer, see above). The result was the binary plasmid vector pMH000-0.

10 The transformation of hypocotyl explants of rape of the variety Drakkar was carried out according to the protocol of De Block (Plant Physiol. 91 (1989), 694-701) by means of Agrobacteria (strain GV 3101 C58C1 Rif^r) carrying the binary vector pMH0208 (ATP/ADP transporter sense). Shoots were regenerated on selective
15 nutrient medium (sulfonamide) and cultivated in the greenhouse up to seed maturation. By means of PCR and leaf test (tolerance against glufosinatammonium (Basta[®])) it was tested which plants contained the transgene. Maturing embryos at various developmental stages were harvested from said plants and stored in liquid nitrogen.

For the determination of the oil content mature seeds of transgenic rape lines and of
20 control lines were analysed by means of the non-invasive near infrared spectroscopy (described, for example, by Schulz et al., J. Near Infrared Spectrosc. 6, (1998), A125-A130; Starr et al., J. Agric. Sci. 104(2), (1985), 317-323).

AMT 34 AMDT

Claims

1. A transgenic plant cell which is genetically modified, wherein the genetic modification is an introduction of a foreign nucleic acid molecule whose presence or expression leads to an increase in the plastidial ADP/ATP translocator activity in comparison with corresponding non-genetically modified plant cells from wild type plants.
2. The transgenic plant cell according to claim 1, wherein the foreign nucleic acid molecule encodes a plastidial ADP/ATP translocator.
3. The transgenic plant cell according to claim 2, wherein the nucleic acid molecule encodes a plastidial ADP/ATP translocator from *Arabidopsis thaliana*.
4. The transgenic plant cell according to any one of claims 1 to 3 exhibiting an increased yield in comparison with corresponding non-genetically modified plant cells.
5. The transgenic plant cell according to any one of claims 1 to 4 exhibiting an increased oil and/or starch content in comparison with corresponding non-genetically modified plant cells.
6. The transgenic plant cell according to any one of claims 1 to 5 synthesizing a starch exhibiting an increased amylose content in comparison with starch from corresponding non-genetically modified plant cells.
7. A transgenic plant containing transgenic plant cells according to any one of claims 1 to 6.
8. The transgenic plant according to claim 7, which is an oil and/or starch storing plant.
9. The transgenic plant according to claim 8, which is a maize, rape, wheat or

potato plant.

10. A method for the production of a transgenic plant exhibiting an increased yield in comparison with wild type plants, wherein
 - (a) a plant cell is genetically modified by means of introduction of a foreign nucleic acid molecule whose presence or expression leads to an increase in the plastidial ADP/ATP translocator activity in the cell;
 - (b) a plant is regenerated from the cell produced according to step (a); and
 - (c) further plants are optionally produced from the plant produced according to step (b).
11. The method according to claim 10, wherein the transgenic plant exhibits an increased oil and/or starch content in comparison with wild type plants and/or whose starch exhibits an increased amylose content in comparison with starch from wild type plants
12. A transgenic plant obtainable by the method according to claim 10 or 11.
13. Propagation material of plants according to any one of claims 7 to 9 or 12, wherein said propagation material contains transgenic cells according to any one of claims 1 to 6.
14. Use of nucleic acid molecules encoding a plastidial ADP/ATP translocator for the production of transgenic plants exhibiting an increased yield in comparison with wild type plants.
15. The use according to claim 14, wherein the transgenic plant exhibits an increased oil and/or starch content and/or synthesizes a starch exhibiting an increased amylose content in comparison with starch from wild type plants.
16. A method for the production of a modified starch comprising the extraction of the starch from a plant according to any one of claims 7 to 9 or according to claim 12.

Abstract

5 Transgenic plant cells and plants are described which, compared to wildtype cells and plants, exhibit an increased yield, in particular, an increased oil and/or starch content and which preferably synthesize a modified starch. The described plants exhibit an increase or a decrease of the plastidial ADP/ATP translocator activity.

1/8




Fig. 1




Fig. 2

DC Alignment:

3/8

1 AATP1_A.T._
 2 AATP2_A.t._
 3 TLC_

```

      10   20   30   40   50   60
1 MEAVIQTRGLLSLPTKPIGVRSQLP SHGLKQRLFAAKPRNLHGCLYPLTGTRNFKPLSQ 60
2 MEGLIQTRGILSLPASHR-SEKVLQPSHGLKQRLFTTN---LPALSLSLMVTRNFKPFSK 56
3 -----MS-----TSK-----SENY--LS- 11

      70   80   90  100  110  120
1 PCMGFRFPTKREAPSSYARRRRGCWRRSCLRRSDSAAVVASRKIFGVEVATLKKIPLGL 120
2 SHLGFRFPTTRREAEDSLARRKLRRPRRCVDEGDTAAMAVSPKIFGVEVTTLKKIVPLGL 116
3 -----EL-----RKIIWP-----IEQYENKKFLPLAF 33

     130   140   150   160   170   180
1 MFFCILFNITLRTDKDVLVVTAKGSSAEIIPFLK TWVNLPMAGFMMLLYTKLSNVLSKK 180
2 MFFCILFNITLRTDKDVLVVTAKGSSAEIIPFLK TWVNLPMAGFMMLLYTKLSNVLSKK 176
3 MMFCILLNYSTLRSIKDGFVVTDIG-T-ESISFLKTYIVLPSAVIAMIIYVKLCDILKQE 91

     190   200   210   220   230   240
1 ALFYTVIVPFIIFYGGFGFVMYPLSNYHPEALADKLLTTLGPRFMGPAILRIWSFCLF 240
2 ALFYTVIVPFIIVYFGAFGFVMYPRSNLIQPEALADKLLATLGPRFMGPALMRIWSFCLF 236
3 NVFYVITSFFLGYFALFAFVLYPYPDLVHPDHKTIESLSLAYPNFKWFIKIVGKWSFASF 151

     250   260   270   280   290   300
1 YVMAELWGSVVSVLFWGFGANQITTVDEAKKFYPLFGIGANVALIFSGRTVKYFSNLRKN 300
2 YVMAELWGSVVSVLFWGFGANQITTVDEAKKFYPLFGLGANVALIFSGRTVKYFSNMRKN 296
3 YTIAELWGTMMLSLLFWQFANQITKIAEAKRFYSMFGLLANLALPVTSSVIGYFLHEKTQ 211

     310   320   330   340   350   360
1 LGPGVDGGSFV---ESHDEHCGGNGTRICLSIGGSNRYV---P-L--PTRSKNKKEKPKMGT 352
2 LGPGVDGWAVSLKAMMSIVVGMGLAICFLYWWVNRYV---P-L--PTRSKKKKVKPQMGT 350
3 I-VAEHLKFV---PLFVIMITSSFLIILT YRWMNKNVLTDPRLYDPALVKEKKTKAKLSF 267

     370   380   390   400   410   420
1 MESLKFLVSSPYIRDLATLVVAYGISINLVEVTWKSCLKAQFPSPNEYSAFMGAFSTCTG 412
2 MESLKFLVSSPYIRDLATLVVAYGISINLVEVTWKSCLKSQFPSPNEYSAFMGDFSTCTG 410
3 IESLKMIFTSKYVGYIALLIHAYGVSVNLVEGVWKS VKELYPTKEAYTIYMGQFQFYQG 327

     430   440   450   460   470   480
1 VATFTMMLLSQYVFNKYGWGVAAKITPTVLLLTGVAFFSLILFGGPFA-PLVAKLGMTPL 471
2 IATFTMMLLSQYVFKKYGWGVAAKITPTVLLLTGVAFFSLILFGGPFA-PLVAKLGMTPL 469
3 WVAIAFMLIGSNILRKVSWLTAAMITPLMMFITGAFFSFIFFDSVIAMNLTGILASSPL 387

     490   500   510   520   530   540
1 LAAVYVGALQNI FSK-SAKYSLFDPCKEMAYIPLDEDTKVKGKAAIDVVCNPLGKSGGAL 530
2 LAAVYVVPPEVSSARVQVQHSSTPSAMQECLYPLDEVSKVKAKLQL-MWSATIGKSGGAL 528
3 TLAVMIGMIQNVL SK-GVKYSLFDATKNMAYIPLDKDLRVKGQA AVEVIGRLGKSGGAI 446

     550   560   570   580   590   600
1 IQQ-FMIL--SFGSLANSTPYLGMILLVIVTAWLAAAKSLEGQFNSLRLKKS LRRKWREL 587
2 IQQ-FMIL--TFGSLANSTPYLG VILLGIVTAWLAAAKSLEGPV----- 569
3 IQSTFFILFPVFGFI-EATPYFASIFFIIVILWIFAVKGLNKEYQVL-VNKNEK----- 498

1 HR 589
2 -- 569
3 -- 498

```

Fig. 3

4/8




Fig. 4

5/8




Fig. 5




Fig. 6

6/8




Fig. 7




Fig. 8




Fig. 9

PLEASE NOTE:
YOU MUST
COMPLETE THE
FOLLOWING



BIRCH, STEWART, KOLASCH & BIRCH, LLP

P.O. Box 747 • Falls Church, Virginia 22040-0747
Telephone: (703) 205-8000 • Facsimile: (703) 205-8050

COMBINED DECLARATION AND POWER OF ATTORNEY FOR PATENT AND DESIGN APPLICATIONS

As a below named inventor, I hereby declare that: my residence, post office address and citizenship are as stated next to my name; that I verily believe that I am the original, first and sole inventor (if only one inventor is named below) or an original, first and joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

Insert Title:

TRANSGENIC PLANTS WITH A MODIFIED ACTIVITY OF A PLASTIDIAL ADP/ATP TRANSLOCATOR

Fill in Appropriate
Information -
For Use Without
Specification
Attached:

the specification of which is attached hereto. If not attached hereto,

the specification was filed on November 6, 2000 as
United States Application Number 09/674,768;
and amended on November 6, 2000 (if applicable) and/or
the specification was filed on May 12, 1999 as PCT
International Application Number PCT/EP99/03292; and was
amended under PCT Article 19 on _____ (if applicable)

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I do not know and do not believe the same was ever known or used in the United States of America before my or our invention thereof, or patented or described in any printed publication in any country before my or our invention thereof or more than one year prior to this application, that the same was not in public use or on sale in the United States of America more than one year prior to this application, that the invention has not been patented or made the subject of an inventor's certificate issued before the date of this application in any country foreign to the United States of America on an application filed by me or my legal representative or assigns more than twelve months (six months for designs) prior to this application, and that no application for patent or inventor's certificate on this invention has been filed in any country foreign to the United States of America prior to this application by me or my legal representatives or assigns, except as follows.

I hereby claim foreign priority benefits under Title 35, United States Code, §119(a)-(d) of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Prior Foreign Application(s)

Priority Claimed

Insert Priority
Information:
(if appropriate)

<u>198 21 442.1</u> (Number)	<u>Germany</u> (Country)	<u>May 13, 1998</u> (Month/Day/Year Filed)	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No
_____ (Number)	_____ (Country)	_____ (Month/Day/Year Filed)	<input type="checkbox"/> Yes	<input type="checkbox"/> No
_____ (Number)	_____ (Country)	_____ (Month/Day/Year Filed)	<input type="checkbox"/> Yes	<input type="checkbox"/> No
_____ (Number)	_____ (Country)	_____ (Month/Day/Year Filed)	<input type="checkbox"/> Yes	<input type="checkbox"/> No

I hereby claim the benefit under Title 35, United States Code, §119(e) of any United States provisional applications(s) listed below.

Insert Provisional
Application(s):
(if any)

_____ (Application Number)	_____ (Filing Date)
_____ (Application Number)	_____ (Filing Date)

All Foreign Applications, if any, for any Patent, or Inventor's Certificate Filed More than 12 Months (6 Months for Designs) Prior to the Filing Date of This Application:

Country	Application Number	Date of Filing (Month/Day/Year)
_____	_____	_____
_____	_____	_____

Insert Requested
Information:
(if appropriate)

I hereby claim the benefit under Title 35, United States Code, §120 of any United States and/or PCT application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States and/or PCT application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose information which is material to the patentability as defined in Title 37, Code of Federal Regulations, §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

Insert Prior U.S.
Application(s):
(if any)

_____ (Application Number)	_____ (Filing Date)	_____ (Status - patented, pending, abandoned)
_____ (Application Number)	_____ (Filing Date)	_____ (Status - patented, pending, abandoned)

I hereby appoint the following attorneys to prosecute this application and/or an international application based on this application and to transact all business in the Patent and Trademark Office connected therewith and in connection with the resulting patent based on instructions received from the entity who first sent the application papers to the attorneys identified below, unless the inventor(s) or assignee provides said attorneys with a written notice to the contrary:

Raymond C. Stewart

(Reg. No. 21,066)

Joseph A. Kolasch

(Reg. No. 22,463)

Bernard L. Sweeney

(Reg. No. 24,448)

Charles Gorenstein

(Reg. No. 29,271)

Leonard R. Svensson

(Reg. No. 30,330)

Andrew D. Meikle

(Reg. No. 32,868)

Joe McKinney Muncy

(Reg. No. 32,334)

John W. Bailey

(Reg. No. 32,881)

Gary D. Yacura

(Reg. No. 35,416)

Terrell C. Birch

(Reg. No. 19,382)

James M. Slattery

(Reg. No. 28,380)

Michael K. Mutter

(Reg. No. 29,680)

Gerald M. Murphy, Jr.

(Reg. No. 28,977)

Terry L. Clark

(Reg. No. 32,644)

Marc S. Weiner

(Reg. No. 32,181)

Donald J. Daley

(Reg. No. 34,313)

John A. Castellano

(Reg. No. 35,094)

O I P E J C I S

FEB 23 2001

PATENT & TRADEMARK OFFICE

Send Correspondence to:

BIRCH, STEWART, KOLASCH & BIRCH, LLPor Customer No. 2292

P.O. Box 747 • Falls Church, Virginia 22040-0747

Telephone: (703) 205-8000 • Facsimile: (703) 205-8050

**PLEASE NOTE:
YOU MUST
COMPLETE
THE
FOLLOWING:**

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full Name of First
or Sole Inventor
Insert Name of
Inventor
Insert Date This
Document is Signed

GIVEN NAME/FAMILY NAME

INVENTOR'S SIGNATURE

DATE*

Ekkehard, NEUHAUS

E. Neuhaus

22.12.2000

Residence (City, State & Country)

CITIZENSHIP

Osnabruck, Germany DEX

German

MAILING ADDRESS (Complete Street Address including City, State & Country)

Sachsenweg 29 49086 Osnabruck, Germany

GIVEN NAME/FAMILY NAME

INVENTOR'S SIGNATURE

DATE*

Torsten, MOEHLMANN

T. Moehlmann

22.12.2000

Residence (City, State & Country)

CITIZENSHIP

Bunde, Germany DEX

German

MAILING ADDRESS (Complete Street Address including City, State & Country)

Gellertstr. 17 32257 Bunde, Germany

GIVEN NAME/FAMILY NAME

INVENTOR'S SIGNATURE

DATE*

Karl-Heinz, GRAEVE-KAMPFENKEL

K. Graeve-Kampfendel

31.12.2000

Residence (City, State & Country)

CITIZENSHIP

Mainz-Kostheim, Germany DEX

German

MAILING ADDRESS (Complete Street Address including City, State & Country)

Im Sempel 10A 55246 Mainz-Kostheim, Germany

GIVEN NAME/FAMILY NAME

INVENTOR'S SIGNATURE

DATE*

Joachim, TJADEN

J. Tjaden

22.12.2000

Residence (City, State & Country)

CITIZENSHIP

Osnabruck, Germany DEX

German

MAILING ADDRESS (Complete Street Address including City, State & Country)

Johannisstr. 97 49074 Osnabruck, Germany

3-00

Full Name of Third
Inventor, if any:
see above

4-00

Full Name of Fourth
Inventor, if any:
see above

SEQUENCE LISTING

<110> NEUHAUS, Ekkehard
 MOEHLMANN, Torsten
 GRAEVE-KAMPFENKEL, Karl-Heinz
 TJADEN, Joachim
 SCHELL, Jozef
 MARTINI, Norbert

<120> Transgenic Plants with a Modified Activity of a Plastidial ADP/ATP
 Translocator

<130> 0147-0215P

<140>

<141>

<150> 198 21 442.1 Germany

<151> 1998-05-13

<150> PCT/EP99/03292

<151> 1999-05-12

<160> 7

<170> PatentIn Ver. 2.1

<210> 1

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: sense-primer

<400> 1

cgtgagagat agagagctcg agggctctgat tcaaacc

37

<210> 2

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: antisense-primer

<400> 2

gatacaacag gaatcctgga tgaagc

26

<210> 3

<211> 56

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 3
gaattcctgc agcccggggg atccactagt ctcgagaagt ggctgggggc ctttcc 56

<210> 4

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 4
tctagaggcc aaggcggccg cttcaacgga ctgcagtgc 39

<210> 5

<211> 589

<212> PRT

<213> Arabidopsis thaliana

<400> 5

Met	Glu	Ala	Val	Ile	Gln	Thr	Arg	Gly	Leu	Leu	Ser	Leu	Pro	Thr	Lys	1	5	10	15
Pro	Ile	Gly	Val	Arg	Ser	Gln	Leu	Gln	Pro	Ser	His	Gly	Leu	Lys	Gln	20	25	30	
Arg	Leu	Phe	Ala	Ala	Lys	Pro	Arg	Asn	Leu	His	Gly	Cys	Leu	Tyr	Pro	35	40	45	
Leu	Thr	Gly	Thr	Arg	Asn	Phe	Lys	Pro	Leu	Ser	Gln	Pro	Cys	Met	Gly	50	55	60	
Phe	Arg	Phe	Pro	Thr	Lys	Arg	Glu	Ala	Pro	Ser	Ser	Tyr	Ala	Arg	Arg	65	70	75	80
Arg	Arg	Gly	Cys	Trp	Arg	Arg	Ser	Cys	Leu	Arg	Arg	Ser	Asp	Ser	Ala	85	90	95	
Ala	Val	Val	Ala	Ser	Arg	Lys	Ile	Phe	Gly	Val	Glu	Val	Ala	Thr	Leu	100	105	110	
Lys	Lys	Ile	Ile	Pro	Leu	Gly	Leu	Met	Phe	Phe	Cys	Ile	Leu	Phe	Asn	115	120	125	
Tyr	Thr	Ile	Leu	Arg	Asp	Thr	Lys	Asp	Val	Leu	Val	Val	Thr	Ala	Lys	130	135	140	
Gly	Ser	Ser	Ala	Glu	Ile	Ile	Pro	Phe	Leu	Lys	Thr	Trp	Val	Asn	Leu	145	150	155	160
Pro	Met	Ala	Ile	Gly	Phe	Met	Leu	Leu	Tyr	Thr	Lys	Leu	Ser	Asn	Val	165	170	175	

Leu Ser Lys Lys Ala Leu Phe Tyr Thr Val Ile Val Pro Phe Ile Ile
 180 185 190
 Tyr Phe Gly Gly Phe Gly Phe Val Met Tyr Pro Leu Ser Asn Tyr Ile
 195 200 205
 His Pro Glu Ala Leu Ala Asp Lys Leu Leu Thr Thr Leu Gly Pro Arg
 210 215 220
 Phe Met Gly Pro Ile Ala Ile Leu Arg Ile Trp Ser Phe Cys Leu Phe
 225 230 235 240
 Tyr Val Met Ala Glu Leu Trp Gly Ser Val Val Val Ser Val Leu Phe
 245 250 255
 Trp Gly Phe Ala Asn Gln Ile Thr Thr Val Asp Glu Ala Lys Lys Phe
 260 265 270
 Tyr Pro Leu Phe Gly Ile Gly Ala Asn Val Ala Leu Ile Phe Ser Gly
 275 280 285
 Arg Thr Val Lys Tyr Phe Ser Asn Leu Arg Lys Asn Leu Gly Pro Gly
 290 295 300
 Val Asp Gly Ser Phe Val Glu Ser His Asp Glu His Cys Gly Gly Asn
 305 310 315 320
 Gly Thr Arg Ile Cys Leu Ser Ile Gly Gly Ser Asn Arg Tyr Val Pro
 325 330 335
 Leu Pro Thr Arg Ser Lys Asn Lys Lys Glu Lys Pro Lys Met Gly Thr
 340 345 350
 Met Glu Ser Leu Lys Phe Leu Val Ser Ser Pro Tyr Ile Arg Asp Leu
 355 360 365
 Ala Thr Leu Val Val Ala Tyr Gly Ile Ser Ile Asn Leu Val Glu Val
 370 375 380
 Thr Trp Lys Ser Lys Leu Lys Ala Gln Phe Pro Ser Pro Asn Glu Tyr
 385 390 395 400
 Ser Ala Phe Met Gly Ala Phe Ser Thr Cys Thr Gly Val Ala Thr Phe
 405 410 415
 Thr Met Met Leu Leu Ser Gln Tyr Val Phe Asn Lys Tyr Gly Trp Gly
 420 425 430
 Val Ala Ala Lys Ile Thr Pro Thr Val Leu Leu Leu Thr Gly Val Ala
 435 440 445
 Phe Phe Ser Leu Ile Leu Phe Gly Gly Pro Phe Ala Pro Leu Val Ala
 450 455 460
 Lys Leu Gly Met Thr Pro Leu Leu Ala Ala Val Tyr Val Gly Ala Leu
 465 470 475 480

Gln Asn Ile Phe Ser Lys Ser Ala Lys Tyr Ser Leu Phe Asp Pro Cys
 485 490 495
 Lys Glu Met Ala Tyr Ile Pro Leu Asp Glu Asp Thr Lys Val Lys Gly
 500 505 510
 Lys Ala Ala Ile Asp Val Val Cys Asn Pro Leu Gly Lys Ser Gly Gly
 515 520 525
 Ala Leu Ile Gln Gln Phe Met Ile Leu Ser Phe Gly Ser Leu Ala Asn
 530 535 540
 Ser Thr Pro Tyr Leu Gly Met Ile Leu Leu Val Ile Val Thr Ala Trp
 545 550 555 560
 Leu Ala Ala Ala Lys Ser Leu Glu Gly Gln Phe Asn Ser Leu Arg Leu
 565 570 575
 Lys Lys Ser Leu Arg Arg Lys Trp Arg Glu Leu His Arg
 580 585

<210> 6
 <211> 569
 <212> PRT
 <213> Arabidopsis thaliana

<400> 6
 Met Glu Gly Leu Ile Gln Thr Arg Gly Ile Leu Ser Leu Pro Ala Ser
 1 5 10 15
 His Arg Ser Glu Lys Val Leu Gln Pro Ser His Gly Leu Lys Gln Arg
 20 25 30
 Leu Phe Thr Thr Asn Leu Pro Ala Leu Ser Leu Ser Leu Met Val Thr
 35 40 45
 Arg Asn Phe Lys Pro Phe Ser Lys Ser His Leu Gly Phe Arg Phe Pro
 50 55 60
 Thr Arg Arg Glu Ala Glu Asp Ser Leu Ala Arg Arg Lys Leu Arg Arg
 65 70 75 80
 Pro Arg Arg Lys Cys Val Asp Glu Gly Asp Thr Ala Ala Met Ala Val
 85 90 95
 Ser Pro Lys Ile Phe Gly Val Glu Val Thr Thr Leu Lys Lys Ile Val
 100 105 110
 Pro Leu Gly Leu Met Phe Phe Cys Ile Leu Phe Asn Tyr Thr Ile Leu
 115 120 125
 Arg Asp Thr Lys Asp Val Leu Val Val Thr Ala Lys Gly Ser Ser Ala
 130 135 140
 Glu Ile Ile Pro Phe Leu Lys Thr Trp Val Asn Val Pro Met Ala Ile

145 150 155 160
 Gly Phe Met Leu Leu Tyr Thr Lys Leu Ser Asn Val Leu Ser Lys Lys
 165 170 175
 Ala Leu Phe Tyr Thr Val Ile Val Pro Phe Ile Val Tyr Phe Gly Ala
 180 185 190
 Phe Gly Phe Val Met Tyr Pro Arg Ser Asn Leu Ile Gln Pro Glu Ala
 195 200 205
 Leu Ala Asp Lys Leu Leu Ala Thr Leu Gly Pro Arg Phe Met Gly Pro
 210 215 220
 Leu Ala Ile Met Arg Ile Trp Ser Phe Cys Leu Phe Tyr Val Met Ala
 225 230 235 240
 Glu Leu Trp Gly Ser Val Val Val Ser Val Leu Phe Trp Gly Phe Ala
 245 250 255
 Asn Gln Ile Thr Thr Val Asp Glu Ala Lys Lys Phe Tyr Pro Leu Phe
 260 265 270
 Gly Leu Gly Ala Asn Val Ala Leu Ile Phe Ser Gly Arg Thr Val Lys
 275 280 285
 Tyr Phe Ser Asn Met Arg Lys Asn Leu Gly Pro Gly Val Asp Gly Trp
 290 295 300
 Ala Val Ser Leu Lys Ala Met Met Ser Ile Val Val Gly Met Gly Leu
 305 310 315 320
 Ala Ile Cys Phe Leu Tyr Trp Trp Val Asn Arg Tyr Val Pro Leu Pro
 325 330 335
 Thr Arg Ser Lys Lys Lys Lys Val Lys Pro Gln Met Gly Thr Met Glu
 340 345 350
 Ser Leu Lys Phe Leu Val Ser Ser Pro Tyr Ile Arg Asp Leu Ala Thr
 355 360 365
 Leu Val Val Ala Tyr Gly Ile Ser Ile Asn Leu Val Glu Val Thr Trp
 370 375 380
 Lys Ser Lys Leu Lys Ser Gln Phe Pro Ser Pro Asn Glu Tyr Ser Ala
 385 390 395 400
 Phe Met Gly Asp Phe Ser Thr Cys Thr Gly Ile Ala Thr Phe Thr Met
 405 410 415
 Met Leu Leu Ser Gln Tyr Val Phe Lys Lys Tyr Gly Trp Gly Val Ala
 420 425 430
 Ala Lys Ile Thr Pro Thr Val Leu Leu Leu Thr Gly Val Ala Phe Phe
 435 440 445
 Ser Leu Ile Leu Phe Gly Gly Pro Phe Ala Pro Leu Val Ala Lys Leu

450 455 460
 Gly Met Thr Pro Leu Leu Ala Ala Val Tyr Val Val Pro Pro Glu Val
 465 470 475 480
 Ser Ser Ala Arg Val Gln Val Gln His Ser Ser Thr Pro Ser Ala Met
 485 490 495
 Gln Glu Cys Leu Tyr Pro Leu Asp Glu Val Ser Lys Val Lys Ala Lys
 500 505 510
 Leu Gln Leu Met Trp Ser Ala Thr Ile Gly Lys Ser Gly Gly Ala Leu
 515 520 525
 Ile Gln Gln Phe Met Ile Leu Thr Phe Gly Ser Leu Ala Asn Ser Thr
 530 535 540
 Pro Tyr Leu Gly Val Ile Leu Leu Gly Ile Val Thr Ala Trp Leu Ala
 545 550 555 560
 Ala Ala Lys Ser Leu Glu Gly Pro Val
 565

<210> 7
 <211> 498
 <212> PRT
 <213> Rickettsia prowazekii

<400> 7
 Met Ser Thr Ser Lys Ser Glu Asn Tyr Leu Ser Glu Leu Arg Lys Ile
 1 5 10 15
 Ile Trp Pro Ile Glu Gln Tyr Glu Asn Lys Lys Phe Leu Pro Leu Ala
 20 25 30
 Phe Met Met Phe Cys Ile Leu Leu Asn Tyr Ser Thr Leu Arg Ser Ile
 35 40 45
 Lys Asp Gly Phe Val Val Thr Asp Ile Gly Thr Glu Ser Ile Ser Phe
 50 55 60
 Leu Lys Thr Tyr Ile Val Leu Pro Ser Ala Val Ile Ala Met Ile Ile
 65 70 75 80
 Tyr Val Lys Leu Cys Asp Ile Leu Lys Gln Glu Asn Val Phe Tyr Val
 85 90 95
 Ile Thr Ser Phe Phe Leu Gly Tyr Phe Ala Leu Phe Ala Phe Val Leu
 100 105 110
 Tyr Pro Tyr Pro Asp Leu Val His Pro Asp His Lys Thr Ile Glu Ser
 115 120 125
 Leu Ser Leu Ala Tyr Pro Asn Phe Lys Trp Phe Ile Lys Ile Val Gly
 130 135 140

Lys Trp Ser Phe Ala Ser Phe Tyr Thr Ile Ala Glu Leu Trp Gly Thr
 145 150 155 160
 Met Met Leu Ser Leu Leu Phe Trp Gln Phe Ala Asn Gln Ile Thr Lys
 165 170 175
 Ile Ala Glu Ala Lys Arg Phe Tyr Ser Met Phe Gly Leu Leu Ala Asn
 180 185 190
 Leu Ala Leu Pro Val Thr Ser Val Val Ile Gly Tyr Phe Leu His Glu
 195 200 205
 Lys Thr Gln Ile Val Ala Glu His Leu Lys Phe Val Pro Leu Phe Val
 210 215 220
 Ile Met Ile Thr Ser Ser Phe Leu Ile Ile Leu Thr Tyr Arg Trp Met
 225 230 235 240
 Asn Lys Asn Val Leu Thr Asp Pro Arg Leu Tyr Asp Pro Ala Leu Val
 245 250 255
 Lys Glu Lys Lys Thr Lys Ala Lys Leu Ser Phe Ile Glu Ser Leu Lys
 260 265 270
 Met Ile Phe Thr Ser Lys Tyr Val Gly Tyr Ile Ala Leu Leu Ile Ile
 275 280 285
 Ala Tyr Gly Val Ser Val Asn Leu Val Glu Gly Val Trp Lys Ser Lys
 290 295 300
 Val Lys Glu Leu Tyr Pro Thr Lys Glu Ala Tyr Thr Ile Tyr Met Gly
 305 310 315 320
 Gln Phe Gln Phe Tyr Gln Gly Trp Val Ala Ile Ala Phe Met Leu Ile
 325 330 335
 Gly Ser Asn Ile Leu Arg Lys Val Ser Trp Leu Thr Ala Ala Met Ile
 340 345 350
 Thr Pro Leu Met Met Phe Ile Thr Gly Ala Ala Phe Phe Ser Phe Ile
 355 360 365
 Phe Phe Asp Ser Val Ile Ala Met Asn Leu Thr Gly Ile Leu Ala Ser
 370 375 380
 Ser Pro Leu Thr Leu Ala Val Met Ile Gly Met Ile Gln Asn Val Leu
 385 390 395 400
 Ser Lys Gly Val Lys Tyr Ser Leu Phe Asp Ala Thr Lys Asn Met Ala
 405 410 415
 Tyr Ile Pro Leu Asp Lys Asp Leu Arg Val Lys Gly Gln Ala Ala Val
 420 425 430
 Glu Val Ile Gly Gly Arg Leu Gly Lys Ser Gly Gly Ala Ile Ile Gln
 435 440 445

